

CD28 Polyclonal Antibody

Rabbit Anti Human Polyclonal Antibody
Catalog # ABV11722

Product Information

Application	WB, E
Primary Accession	P10747
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	25066

Additional Information

Gene ID	940
Positive Control	WB, ELISA
Application & Usage	Western Blot: 1:500 – 1:2000, ELISA: 1:20000.
Other Names	CD28; T-cell-specific surface glycoprotein CD28; TP44; CD28
Target/Specificity	CD28
Antibody Form	Liquid
Appearance	Colorless liquid
Formulation	PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Handling	The antibody solution should be gently mixed before use.
Reconstitution & Storage	-20 °C
Background Descriptions	
Precautions	CD28 Polyclonal Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CD28
Function	Receptor that plays a role in T-cell activation, proliferation, survival and the maintenance of immune homeostasis (PubMed: 1650475 , PubMed: 7568038). Functions not only as an amplifier of TCR signals but delivers unique signals that control intracellular biochemical events that alter the gene expression program of T-cells (PubMed: 24665965). Stimulation upon engagement of its

cognate ligands CD80 or CD86 increases proliferation and expression of various cytokines in particular IL2 production in both CD4(+) and CD8(+) T-cell subsets (PubMed:[12196291](#), PubMed:[1650475](#), PubMed:[35397202](#)). Mechanistically, ligation induces recruitment of protein kinase C-theta/PRKCQ and GRB2 leading to NF-kappa-B activation via both PI3K/Akt-dependent and -independent pathways (PubMed:[21964608](#), PubMed:[24665965](#), PubMed:[7568038](#)). In conjunction with TCR/CD3 ligation and CD40L costimulation, enhances the production of IL4 and IL10 in T- cells (PubMed:[8617933](#)).

Cellular Location

Cell membrane; Single-pass type I membrane protein


Tissue Location

Expressed in T-cells and plasma cells, but not in less mature B-cells

Background

Involved in T-cell activation, induction of cell proliferation and cytokine production and promotion of T-cell survival. Enhances the production of IL4 and IL10 in T-cells in conjunction with TCR/CD3 ligation and CD40L costimulation. Isoform 3 enhances CD40L-mediated activation of NF-kappa-B and kinases MAPK8 and PAK2 in T-cells.

Images



Western blot analysis of K562 cells using CD28 Polyclonal antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.