

# Phospho-ANAPC1 (Ser688) Antibody

Rabbit Polyclonal Antibody Catalog # ABV11850

#### **Product Information**

**Application** WB, IHC, IF, ICC

Primary Accession Q9H1A4

**Reactivity** Human, Mouse

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Calculated MW 216500

#### **Additional Information**

**Gene ID** 64682

Positive Control WB: HEK293; IHC: human breast cancer tissue; IFC: HEK293 cells

**Application & Usage** WB; 1:500 – 1:2000, IHC; 1:50 – 1:200, IF/IC; 1:50 – 1:100

Alias Symbol ANAPC1

Other Names TSG24, Anaphase-promoting complex subunit 1, APC1, Cyclosome subunit 1,

Mitotic checkpoint regulator, Testis-specific gene 24 protein

**Appearance** Colorless liquid

**Formulation** In 0.42% Potassium phosphate; 0.87% Sodium chloride; pH 7.3; 30% glycerol

and 0.01% sodium azide

Reconstitution & Storage -20 °C

**Background Descriptions** 

**Precautions** Phospho-ANAPC1 (Ser688) Antibody is for research use only and not for use

in diagnostic or therapeutic procedures.

#### **Protein Information**

Name ANAPC1

Synonyms TSG24

**Function** Component of the anaphase promoting complex/cyclosome (APC/C), a cell


cycle-regulated E3 ubiquitin ligase that controls progression through mitosis and the G1 phase of the cell cycle (PubMed:<u>18485873</u>). The APC/C complex acts by mediating ubiquitination and subsequent degradation of target proteins: it mainly mediates the formation of 'Lys-11'-linked polyubiquitin chains and, to a lower extent, the formation of 'Lys-48'- and 'Lys-63'-linked

polyubiquitin chains (PubMed:<u>18485873</u>). The APC/C complex catalyzes assembly of branched 'Lys-11'-/'Lys-48'-linked branched ubiquitin chains on target proteins (PubMed:<u>29033132</u>).


## **Background**

Component of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated E3 ubiquitin ligase that controls progression through mitosis and the G1 phase of the cell cycle. The APC/C complex acts by mediating ubiquitination and subsequent degradation of target proteins: it mainly mediates the formation of Lys-11-linked polyubiquitin chains and, to a lower extent, the formation of Lys-48- and Lys-63-linked polyubiquitin chains.


### **Images**



Immunohistochemical analysis of ANAPC1 (pS688) staining in H.breast cancer formalin fixed paraffin embedded tissue section.



Immunofluorescent analysis of ANAPC1(pS688) staining in HEK293T cells.



Western blot analysis of ANAPC1(pS688) expression in HEK293 LPS-treated (A) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.