

LAG3抗体试剂(免疫组织化学)

Rabbit Monoclonal antibody(Mab)

Catalog # AD80600

Product Information

Application	IHC-P
Primary Accession	P18627
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal
Clone Names	412B5F6
Calculated MW	57449

Additional Information

Gene ID	3902
Other Names	Lymphocyte activation gene 3 protein, LAG-3, CD223, Secreted lymphocyte activation gene 3 protein, sLAG-3, LAG3 (HGNC:6476), FDC
Dilution	IHC-P~N/A
Storage	Maintain refrigerated at 2-8°C.

Protein Information

Name	LAG3 {ECO:0000303 PubMed:35761082, ECO:0000312 HGNC:HGNC:6476}
Function	[Lymphocyte activation gene 3 protein]: Inhibitory receptor on antigen activated T-cells (PubMed: 20421648 , PubMed: 35761082 , PubMed: 7805750 , PubMed: 8647185). Delivers inhibitory signals upon binding to ligands, such as MHC class II, its main ligand present at the surface of antigen-presenting cells (APCs), and FGL1, which is secreted by hepatocytes and certain types of tumor cells (PubMed: 30580966 , PubMed: 32920841 , PubMed: 35761082 , PubMed: 39671469 , PubMed: 7589152 , PubMed: 8647185 , PubMed: 9159144). Ligand-binding initiates a signaling that inhibits the T-cell receptor (TCR) in the immunological synapse, preventing T-cell activation (PubMed: 40101708). Mechanistically, ligand-binding promotes (1) ubiquitination of the KIEELE motif, unleashing the RRFSALE motif from the membrane and (2) leading to the formation of condensates with the TCR component CD3E, thereby disrupting the association between CD3E and LCK and preventing TCR activation (PubMed: 40101708 , PubMed: 40592325). May inhibit antigen-specific T-cell activation in synergy with PDCD1/PD-1 (By similarity). Negatively regulates the proliferation, activation, effector function and homeostasis of both CD8(+) and CD4(+) T-cells (PubMed: 20421648 , PubMed: 7805750 , PubMed: 8647185). Also mediates immune tolerance: constitutively expressed on a subset of regulatory T-cells (Tregs) and contributes to their suppressive

Cellular Location	function (By similarity). Also acts as a negative regulator of plasmacytoid dendritic cell (pDCs) activation (By similarity). [Lymphocyte activation gene 3 protein]: Cell membrane; Single-pass type I membrane protein. Note=Clusters on the T-cell surface following ligand-binding
Tissue Location	Primarily expressed in activated T-cells and a subset of natural killer (NK) cells.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.