

# Anti- RACGAP1 / MgcRacGAP Antibody (C-term), Biotinylated

Catalog # AF4273a

## Product Information

---

<b>Application</b>	WB, IHC, E
<b>Primary Accession</b>	<a href="#">Q9H0H5</a>
<b>Other Accession</b>	<a href="#">29127</a> , <a href="#">NP_037409.2</a> , <a href="#">NP_001306934.1</a> , <a href="#">NP_001306935.1</a> , <a href="#">NP_001306936.1</a>
<b>Reactivity</b>	Human
<b>Predicted</b>	Human, Dog
<b>Calculated MW</b>	71027

## Additional Information

---

<b>Gene ID</b>	29127
<b>Other Names</b>	GAP; Rho signaling; spermatogenesis; mitosis; phosphoprotein; acetylprotein; spindlebody
<b>Target/Specificity</b>	This antibody is expected to recognize all reported isoforms (NP_037409.2; NP_001306934.1; NP_001306935.1; NP_001306936.1). Reported variants represent identical protein: NP_001306928.1, NP_001119576.1, NP_037409.2, NP_001306931.1, NP_001306929.1, NP_0011
<b>Dilution</b>	WB~~1:1000 IHC~~1:100~500 E~~N/A
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	Anti- RACGAP1 / MgcRacGAP Antibody (C-term), Biotinylated is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	RACGAP1 ( <a href="#">HGNC:9804</a> )
<b>Function</b>	Component of the centralspindlin complex that serves as a microtubule-dependent and Rho-mediated signaling required for the myosin contractile ring formation during the cell cycle cytokinesis. Required for proper attachment of the midbody to the cell membrane during cytokinesis. Sequentially binds to ECT2 and RAB11FIP3 which regulates cleavage furrow ingression and abscission during cytokinesis (PubMed: <a href="#">18511905</a> ). Plays key roles in controlling cell growth and differentiation of hematopoietic cells through mechanisms other than regulating Rac GTPase activity (PubMed: <a href="#">10979956</a> ). Has a critical role in erythropoiesis (PubMed: <a href="#">34818416</a> ).

Also involved in the regulation of growth-related processes in adipocytes and myoblasts. May be involved in regulating spermatogenesis and in the RACGAP1 pathway in neuronal proliferation. Shows strong GAP (GTPase activation) activity towards CDC42 and RAC1 and less towards RHOA. Essential for the early stages of embryogenesis. May play a role in regulating cortical activity through RHOA during cytokinesis. May participate in the regulation of sulfate transport in male germ cells.

### Cellular Location

Nucleus. Cytoplasm. Cytoplasm, cytoskeleton, spindle Cytoplasmic vesicle, secretory vesicle, acrosome. Cleavage furrow Midbody, Midbody ring. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Midbody. Note=Colocalizes with RND2 in Golgi- derived proacrosomal vesicles and the acrosome (By similarity). During interphase, localized to the nucleus and cytoplasm along with microtubules, in anaphase, is redistributed to the central spindle and, in telophase and cytokinesis, to the midbody ring, also called Flemming body. Colocalizes with RHOA at the myosin contractile ring during cytokinesis. Colocalizes with ECT2 to the mitotic spindles during anaphase/metaphase, the cleavage furrow during telophase and at the midbody at the end of cytokinesis. Colocalizes with Cdc42 to spindle microtubules from prometaphase to telophase. Localizes to the midbody in a KIF23 and PRC1-dependent manner. {ECO:0000250 | UniProtKB:Q9WVM1, ECO:0000269 | PubMed:36825683}

### Tissue Location

Highly expressed in testis, thymus and placenta. Expressed at lower levels in spleen and peripheral blood lymphocytes In testis, expression is restricted to germ cells with the highest levels of expression found in spermatocytes. Expression is regulated in a cell cycle-dependent manner and peaks during G2/M phase

### Images



Biotinylated Antibody (1 µg/ml) staining of Jurkat lysate (35 µg protein in RIPA buffer), exactly mirroring its parental non-biotinylated product. Primary incubation was 1 hour. Detected by chemiluminescence, using streptavidin-HRP and using NAP blocker as a substitute for skimmed milk.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.