

# Goat anti-LIMK2 (aa181-193) Antibody

Peptide-affinity purified goat antibody

Catalog # AF4448a

## Product Information

---

Application	WB, Pep-ELISA
Primary Accession	<a href="#">P53671</a>
Other Accession	<a href="#">NP_001026971.1</a> , <a href="#">NP_005560.1</a> , <a href="#">NP_057952.1</a>
Reactivity	Human, Mouse, Rat, Pig
Host	Goat
Clonality	Polyclonal
Clone Names	LIMK2
Calculated MW	72232

## Additional Information

---

Gene ID	3985
Other Names	LIMK2; LIM domain kinase 2
Dilution	WB~~1:1000 Pep-ELISA~~N/A
Format	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Immunogen	This antibody is expected to recognize all reported isoforms (NP_001026971.1; NP_005560.1; NP_057952.1).
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Goat anti-LIMK2 (aa181-193) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	LIMK2
Function	Serine/threonine-protein kinase that plays an essential role in the regulation of actin filament dynamics (PubMed: <a href="#">10436159</a> , PubMed: <a href="#">11018042</a> ). Acts downstream of several Rho family GTPase signal transduction pathways (PubMed: <a href="#">10436159</a> , PubMed: <a href="#">11018042</a> ). Involved in astral microtubule organization and mitotic spindle orientation during early stages of mitosis by mediating phosphorylation of TPPP (PubMed: <a href="#">22328514</a> ). Displays serine/threonine-specific phosphorylation of myelin basic protein and histone

(MBP) in vitro (PubMed:[8537403](#)). Suppresses ciliogenesis via multiple pathways; phosphorylation of CFL1, suppression of directional trafficking of ciliary vesicles to the ciliary base, and by facilitating YAP1 nuclear localization where it acts as a transcriptional corepressor of the TEAD4 target genes AURKA and PLK1 (PubMed:[25849865](#)).

**Cellular Location**

Cytoplasm, cytoskeleton, spindle. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome [Isoform LIMK2b]: Cytoplasm. Cytoplasm, perinuclear region. Nucleus Note=Mainly present in the cytoplasm and is scarcely translocated to the nucleus.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.