

## Goat anti-ERCC1 Antibody

Peptide-affinity purified goat antibody Catalog # AF4504a

## **Product Information**

Application IF, Pep-ELISA Primary Accession P07992

Other Accession NP 973730.1, NP 001974.1, NP 001159521.1

**Reactivity** Human, Mouse, Rat, Dog, Bovine

HostGoatClonalityPolyclonalClone NamesERCC1Calculated MW32562

## **Additional Information**

Gene ID 2067

**Other Names** ERCC1; excision repair cross-complementing rodent repair deficiency,

complementation group 1 (includes overlapping antisense sequence); COFS4;

UV20; excision repair cross-complementing 1; excision repair protein

**Dilution** IF~~1:50~200 Pep-ELISA~~N/A

**Format** Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5%

bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and

thawing.

**Storage** Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** Goat anti-ERCC1 Antibody is for research use only and not for use in

diagnostic or therapeutic procedures.

## **Protein Information**

Name ERCC1

**Function** [Isoform 1]: Non-catalytic component of a structure-specific DNA repair

endonuclease responsible for the 5'-incision during DNA repair. Responsible,

in conjunction with SLX4, for the first step in the repair of interstrand

cross-links (ICL). Participates in the processing of anaphase bridge-generating DNA structures, which consist in incompletely processed DNA lesions arising during S or G2 phase, and can result in cytokinesis failure. Also required for homology-directed repair (HDR) of DNA double-strand breaks, in conjunction

with SLX4.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.