

Goat anti-Ku70 / G22P1 Antibody

Peptide-affinity purified goat antibody

Catalog # AF4528a

Product Information

Application	FC, Pep-ELISA
Primary Accession	P12956
Other Accession	NP_001460.1 , NP_001275907.1
Reactivity	Human, Mouse, Rat, Dog
Host	Goat
Clonality	Polyclonal
Clone Names	XRCC6
Calculated MW	69843

Additional Information

Gene ID	2547
Other Names	G22P1; thyroid autoantigen 70kDa (Ku antigen); HGNC:4055; CTC75; CTCBF; KU70; ML8; TLAA; ATP-dependent DNA helicase II, 70 kDa subunit; CTC box binding factor 75 kDa subunit; OTTHUMP00000028581; thyroid autoantigen 70kD (Ku antigen); thyroid-lupus autoant
Dilution	FC~1:10~50 Pep-ELISA~~N/A
Format	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Goat anti-Ku70 / G22P1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	XRCC6 (HGNC:4055)
Synonyms	G22P1
Function	DNA-binding protein critical for the DNA damage response, specifically in repairing double-strand breaks (DSBs) via the classical non-homologous end joining (NHEJ) pathway. It forms a heterodimer with XRCC5 (Ku80), creating the Ku70:Ku80 heterodimer (Ku complex), which serves as a DNA end-binding complex. It primarily binds DSBs and recruits essential repair factors,

assembling the core long-range NHEJ complex to facilitate the alignment and ligation of broken DNA ends (PubMed:[11493912](#), PubMed:[20493174](#), PubMed:[33854234](#), PubMed:[34352203](#), PubMed:[9742108](#)). This pathway ensures the rapid repair of cytotoxic and mutagenic DSBs and contributes to the generation of diversity in T-cell receptors and antibodies through mechanisms such as V(D)J recombination (PubMed:[9742108](#)). Likely acts as a 5'-deoxyribose-5-phosphate lyase (5'-dRP lyase), catalyzing the beta-elimination of the 5'-deoxyribose- 5-phosphate at abasic sites near DSBs. This activity cleans the termini of abasic sites, a common form of nucleotide damage, preparing broken ends for ligation (PubMed:[20383123](#)). It may also possess 3'-5' DNA helicase activity, although this has not been confirmed in vivo, and its physiological significance remains unclear (PubMed:[7957065](#)). Beyond DNA repair, the protein contributes to telomere maintenance (PubMed:[29490055](#)). It is also implicated in transcriptional regulation, acting as a cofactor for various transcription factors (PubMed:[12145306](#), PubMed:[8621488](#)). It plays a role in the regulation of DNA virus-mediated innate immune response by assembling into the HDP- RNP complex, a complex that serves as a platform for IRF3 phosphorylation and subsequent innate immune response activation through the cGAS-STING pathway (PubMed:[28712728](#)). Can also bind RNAs and recruits PRKDC to a wide range of cellular RNAs, including the U3 small nucleolar RNA, playing a role in the biogenesis of ribosomal RNAs (PubMed:[32103174](#)). Additionally, it negatively regulates apoptosis by interacting with BAX, sequestering it from the mitochondria, and may possess deubiquitination activity targeting BAX (PubMed:[15023334](#), PubMed:[18362350](#), PubMed:[35545041](#)).

Cellular Location

Nucleus. Chromosome. Cytoplasm. Note=When trimethylated, localizes in the cytoplasm.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.