

# TGF-alpha (Transforming Growth Factor alpha) Antibody - With BSA and Azide

Mouse Monoclonal Antibody [Clone SPM357 ]  
Catalog # AH10761

## Product Information

---

<b>Application</b>	IF, FC
<b>Primary Accession</b>	<a href="#">P01135</a>
<b>Other Accession</b>	<a href="#">7039</a> , <a href="#">170009</a>
<b>Reactivity</b>	Human, Mouse, Rat, Bovine
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	Mouse / IgG1, kappa
<b>Clone Names</b>	SPM357
<b>Calculated MW</b>	6 KDa

## Additional Information

---

<b>Other Names</b>	Protransforming growth factor alpha, Transforming growth factor alpha, TGF-alpha, EGF-like TGF, ETGF, TGF type 1, TGFA
<b>Application Note</b>	IF~~1:50~200 FC~~1:10~50
<b>Format</b>	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage</b>	Store at 2 to 8°C. Antibody is stable for 24 months.
<b>Precautions</b>	TGF-alpha (Transforming Growth Factor alpha) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

### Background

---

This antibody reacts with the C-terminus of TGF alpha and shows no cross-reaction with EGF and the neuropeptide synenkephalin. The staining with Ab-1 is completely blocked by the peptide used for raising this antibody. TGF  $\alpha$  (aa50) is a growth factor with 33% homology to EGF, binds to EGFR, activates tyrosine phosphorylation of the receptor, and stimulates cell proliferation. It plays a role in tumor initiation by inducing the reversible transformed phenotype.

## References

---

Kobrin MS et. al. J Bio Chem, 1986, 261:14414-9. | Kudlow JE et. al. Endocrinology, 1987, 121(4):1577-9. |  
Kobrin MS et. al. Endocrinology, 1987, 121(4):1412-6. | Kudlow JE et. al. J Bio Chem, 1989, 264(7):3880-3 |

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.