



EGFR (Epidermal Growth Factor Receptor) Antibody - With BSA and Azide

Mouse Monoclonal Antibody [Clone SPM622] Catalog # AH11177

Product Information

Application IHC, IF, FC
Primary Accession P00533
Other Accession 1956, 488293
Reactivity Human
Host Mouse
Clonality Monoclonal

Isotype Mouse / IgG1, kappa

Clone Names SPM622 Calculated MW 134277

Additional Information

Gene ID 1956

Other Names Epidermal growth factor receptor, 2.7.10.1, Proto-oncogene c-ErbB-1,

Receptor tyrosine-protein kinase erbB-1, EGFR, ERBB, ERBB1, HER1

Application Note IHC~~1:100~500 IF~~1:50~200 FC~~1:10~50

Storage Store at 2 to 8°C.Antibody is stable for 24 months.

Precautions EGFR (Epidermal Growth Factor Receptor) Antibody - With BSA and Azide is

for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name EGFR (HGNC:3236)

Synonyms ERBB, ERBB1, HER1

Function Receptor tyrosine kinase binding ligands of the EGF family and activating

several signaling cascades to convert extracellular cues into appropriate cellular responses (PubMed:10805725, PubMed:27153536, PubMed:2790960, PubMed:35538033). Known ligands include EGF, TGFA/TGF- alpha, AREG, epigen/EPGN, BTC/betacellulin, epiregulin/EREG and HBEGF/heparin-binding

EGF (PubMed: 12297049, PubMed: 15611079, PubMed: 17909029,

PubMed: 20837704, PubMed: 27153536, PubMed: 2790960, PubMed: 7679104, PubMed: 8144591, PubMed: 9419975). Ligand binding triggers receptor homoand/or heterodimerization and autophosphorylation on key cytoplasmic residues. The phosphorylated receptor recruits adapter proteins like GRB2

which in turn activates complex downstream signaling cascades. Activates at least 4 major downstream signaling cascades including the RAS-RAF-MEK-ERK, PI3 kinase-AKT, PLCgamma-PKC and STATs modules (PubMed:27153536). May also activate the NF-kappa-B signaling cascade (PubMed:11116146). Also directly phosphorylates other proteins like RGS16, activating its GTPase activity and probably coupling the EGF receptor signaling to the G protein-coupled receptor signaling (PubMed:11602604). Also phosphorylates MUC1 and increases its interaction with SRC and CTNNB1/beta-catenin (PubMed:11483589). Positively regulates cell migration via interaction with CCDC88A/GIV which retains EGFR at the cell membrane following ligand stimulation, promoting EGFR signaling which triggers cell migration (PubMed:20462955). Plays a role in enhancing learning and memory performance (By similarity). Plays a role in mammalian pain signaling (long-lasting hypersensitivity) (By similarity).

Cellular Location

Cell membrane; Single-pass type I membrane protein. Endoplasmic reticulum membrane; Single-pass type I membrane protein Golgi apparatus membrane; Single-pass type I membrane protein. Nucleus membrane; Single-pass type I membrane protein. Endosome. Endosome membrane. Nucleus. Note=In response to EGF, translocated from the cell membrane to the nucleus via Golgi and ER (PubMed:17909029, PubMed:20674546). Endocytosed upon activation by ligand (PubMed:17182860, PubMed:17909029, PubMed:27153536, PubMed:2790960). Colocalized with GPER1 in the nucleus of estrogen agonist-induced cancer-associated fibroblasts (CAF) (PubMed:20551055)

Tissue Location

Ubiquitously expressed. Isoform 2 is also expressed in ovarian cancers.


Background

This MAb recognizes a protein of 170kDa, identified as EGFR. EGFR is type I receptor tyrosine kinase with sequence homology to erbB-1, -2, -3 -4 or HER-1, -2, -3 -4. It binds to Epidermal Growth Factor (EGF), Transforming Growth Factor-a (TGF-a), Heparin-binding EGF (HB-EGF), amphiregulin, βcellulin and epiregulin. EGFR is overexpressed in tumors of breast, brain, bladder, lung, gastric, head & neck, esophagus, cervix, vulva, ovary, and endometrium. It is predominantly present in squamous cell carcinomas.

References

Tungekar MF et. al. Journal of Clinical Pathology. 51: 583-587 (1998).

Images



Formalin-fixed, paraffin-embedded human Placenta stained with EGFR Monoclonal Antibody (SPM622).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.