

# CD3e (T-Cell Marker) Antibody - With BSA and Azide

Mouse Monoclonal Antibody [Clone CRIS-7 ]

Catalog # AH12595

## Product Information

---

<b>Application</b>	IF, FC
<b>Primary Accession</b>	<a href="#">P07766</a>
<b>Other Accession</b>	<a href="#">916_3003</a>
<b>Reactivity</b>	Human, Monkey
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	Mouse / IgG2a, kappa
<b>Clone Names</b>	CRIS-7
<b>Calculated MW</b>	23147

## Additional Information

---

<b>Gene ID</b>	916
<b>Other Names</b>	T-cell surface glycoprotein CD3 epsilon chain, T-cell surface antigen T3/Leu-4 epsilon chain, CD3e, CD3E, T3E
<b>Application Note</b>	IF~1:50~200 FC~1:10~50
<b>Storage</b>	Store at 2 to 8°C. Antibody is stable for 24 months.
<b>Precautions</b>	CD3e (T-Cell Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	CD3E
<b>Synonyms</b>	T3E
<b>Function</b>	Part of the TCR-CD3 complex present on T-lymphocyte cell surface that plays an essential role in adaptive immune response (PubMed: <a href="#">15294938</a> , PubMed: <a href="#">15546002</a> , PubMed: <a href="#">2470098</a> , PubMed: <a href="#">40592325</a> , PubMed: <a href="#">8490660</a> ). When antigen presenting cells (APCs) activate T-cell receptor (TCR), TCR-mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD247/CD3Z (PubMed: <a href="#">2470098</a> , PubMed: <a href="#">40592325</a> ). All CD3 chains contain immunoreceptor tyrosine-based activation motifs (ITAMs) in their cytoplasmic domain (PubMed: <a href="#">2470098</a> , PubMed: <a href="#">40592325</a> ). Upon TCR engagement, these motifs become phosphorylated by Src family protein tyrosine kinases LCK and FYN, resulting in the activation of downstream signaling pathways (PubMed: <a href="#">2470098</a> ,

PubMed:[40592325](#)). CD3E ITAM phosphorylation creates docking sites for the protein kinase ZAP70 leading to ZAP70 phosphorylation and its conversion into a catalytically active enzyme (By similarity). In addition of this role of signal transduction in T-cell activation, CD3E plays an essential role in correct T-cell development (By similarity). Also participates in internalization and cell surface down-regulation of TCR-CD3 complexes via endocytosis sequences present in CD3E cytosolic region (PubMed:[10384095](#), PubMed:[26507128](#)). In addition to its role as a TCR coreceptor, it serves as a receptor for ITPR1PL1 (PubMed:[38614099](#)). Ligand recognition inhibits T-cell activation by promoting interaction with NCK1, which prevents CD3E-ZAP70 interaction and blocks the ERK- NFkB signaling cascade and calcium influx (PubMed:[12110186](#), PubMed:[38614099](#)).

<b>Cellular Location</b>	Cell membrane; Single-pass type I membrane protein
--------------------------	--

## Background

---

Recognizes the -chain of CD3 (Workshop V; Code: CD03.09), which consists of five different polypeptide chains (designated as  $\gamma$ ,  $\delta$ ,  $\epsilon$ ,  $\zeta$ , and  $\eta$ ) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage. In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms.

## References

---

McMichael AJ et al. (eds) Leukocyte Typing III, Oxford University Press, Oxford, 1987. Knapp W et al. (eds) Leukocyte Typing IV, p245 and 1059, Oxford University Press, Oxford, 1989 | Schlossman S et al. (eds) Leukocyte Typing V. Oxford University Press, Oxford, 1995. | Alberola-Ila J et al. Stimulation through the TCR/CD3 complex up-regulates the CD2 surface expression on human T lymphocytes. J Immunol 1991, 146(4):108

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.