

# Anti-P-Cadherin (CDH3) Antibody

Mouse Monoclonal Antibody

Catalog # AH13077

## Product Information

---

<b>Application</b>	WB, IF, FC
<b>Primary Accession</b>	<a href="#">P22223</a>
<b>Other Accession</b>	<a href="#">191842</a>
<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	Mouse / IgG1, kappa
<b>Clone Names</b>	6A9
<b>Calculated MW</b>	91418

## Additional Information

---

<b>Gene ID</b>	1001
<b>Other Names</b>	Cadherin 3 type 1; Cadherin-3; Cadp; Calcium dependent adhesion protein placental; CDH3; CDHP; HJMD; P-cadherin (Placental); PCAD; Placental cadherin
<b>Application Note</b>	Flow Cytometry (0.5-1ug/million cells); ,Immunofluorescence (1-2ug/ml); ,Western Blotting (0.5-1.0ug/ml) ,Optimal dilution for a specific application should be determined.
<b>Format</b>	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage</b>	Store at 2 to 8°C.Antibody is stable for 24 months.
<b>Precautions</b>	Anti-P-Cadherin (CDH3) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	CDH3
<b>Synonyms</b>	CDHP
<b>Function</b>	Cadherins are calcium-dependent cell adhesion proteins. They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types.

<b>Cellular Location</b>	Cell membrane; Single-pass type I membrane protein
<b>Tissue Location</b>	Expressed in some normal epithelial tissues and in some carcinoma cell lines.

## **Background**

---

Recognizes a protein of 116kDa, identified as P-Cadherin-1 (CDH3). It is a calcium-dependent cell-cell adhesion glycoprotein comprised of five extracellular cadherin repeats, a transmembrane region and a highly conserved cytoplasmic tail. This gene is located in a six-cadherin cluster in a region on the long arm of chromosome 16 that is involved in loss of heterozygosity events in breast and prostate cancer. In addition, aberrant expression of this protein is observed in cervical adenocarcinomas. Mutations in this gene have been associated with congenital hypotrichosis with juvenile macular dystrophy.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.