

# UHRF2 antibody - middle region

Rabbit Polyclonal Antibody Catalog # AI10140

#### **Product Information**

Application WB
Primary Accession Q96PU4

Other Accession <u>Q96PU4, NP 690856, NM 152896</u>

**Reactivity**Human, Mouse, Rat, Rabbit, Zebrafish, Dog, Guinea Pig, Horse, Bovine **Predicted**Human, Mouse, Rat, Rabbit, Chicken, Dog, Guinea Pig, Horse, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 89985

### **Additional Information**

**Gene ID** 115426

**Alias Symbol** DKFZp434B0920, DKFZp686G0837, MGC33463, NIRF, RNF107, RP11-472F14.2,

URF2

Other Names E3 ubiquitin-protein ligase UHRF2, 632-, Np95/ICBP90-like RING finger

protein, Np95-like RING finger protein, Nuclear protein 97, Nuclear zinc finger protein Np97, RING finger protein 107, Ubiquitin-like PHD and RING finger domain-containing protein 2, Ubiquitin-like-containing PHD and RING finger

domains protein 2, UHRF2, NIRF, RNF107

**Target/Specificity** UHRF2 encodes a nuclear protein which is involved in cell-cycle regulation.

The encoded protein is a ubiquitin-ligase capable of ubiquinating PCNP (PEST-containing nuclear protein), and together they may play a role in tumorigenesis. This gene encodes a nuclear protein which is involved in cell-cycle regulation. The encoded protein is a ubiquitin-ligase capable of ubiquinating PCNP (PEST-containing nuclear protein), and together they may

play a role in tumorigenesis.

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

**Reconstitution & Storage** Add 50 ul of distilled water. Final anti-UHRF2 antibody concentration is 1

mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at

-20°C. Avoid repeat freeze-thaw cycles.

**Precautions** UHRF2 antibody - middle region is for research use only and not for use in

diagnostic or therapeutic procedures.

### **Protein Information**

Name UHRF2

Synonyms NIRF, RNF107

**Function** E3 ubiquitin ligase that plays important roles in DNA methylation, histone

modifications, cell cycle and DNA repair (PubMed:15178429,

PubMed:<u>23404503</u>, PubMed:<u>27743347</u>, PubMed:<u>29506131</u>). Acts as a specific reader for 5-hydroxymethylcytosine (5hmC) and thereby recruits various

substrates to these sites to ubiquitinate them (PubMed:24813944,

PubMed:<u>27129234</u>). This activity also allows the maintenance of 5mC levels at specific genomic loci and regulates neuron-related gene expression (By similarity). Participates in cell cycle regulation by ubiquitinating cyclins CCND1

and CCNE1 and thereby inducing G1 arrest (PubMed: 15178429,

PubMed: 15361834, PubMed: 21952639). Also ubiquitinates PCNP leading to its degradation by the proteasome (PubMed: 12176013, PubMed: 14741369). Plays an active role in DNA damage repair by ubiquitinating p21/CDKN1A leading to its proteasomal degradation (PubMed: 29923055). Also promotes

DNA repair by acting as an interstrand cross-links (ICLs) sensor.

Mechanistically, cooperates with UHRF1 to ensure recruitment of FANCD2 to ICLs, leading to FANCD2 monoubiquitination and subsequent activation (PubMed:30335751). Contributes to UV-induced DNA damage response by physically interacting with ATR in response to irradiation, thereby promoting

ATR activation (PubMed:33848395).

**Cellular Location** Nucleus {ECO:0000255 | PROSITE-ProRule:PRU00358.


ECO:0000269 | PubMed:12176013, ECO:0000269 | PubMed:23404503, ECO:0000269 | PubMed:27129234, ECO:0000269 | PubMed:27743347, ECO:0000269 | PubMed:29923055, ECO:0000269 | PubMed:30335751}. Chromosome. Note=Enriched at genomic loci that are enriched for

5-hydroxymethylcytosine (5hmC)

## **Background**

This is a rabbit polyclonal antibody against UHRF2. It was validated on Western Blot using a cell lysate as a positive control. Abgent strives to provide antibodies covering each member of a whole protein family of your interest. We also use our best efforts to provide you antibodies recognize various epitopes of a target protein. For availability of antibody needed for your experiment, please inquire (sales@abgent.com).

## **Images**



UHRF2 antibody - middle region (AI10140) in Human Liver cells using Western Blot

WB Suggested Anti-UHRF2 Antibody Titration: 0.2-1 μg/ml

ELISA Titer: 1:62500

Positive Control: Human Liver

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.