

# DBP antibody - middle region

Rabbit Polyclonal Antibody Catalog # AI10247

#### **Product Information**

Application WB, IHC Primary Accession Q10586

Other Accession NM 001352, NP 001343

**Reactivity**Human, Mouse, Rat, Pig, Dog, Horse, Bovine, Sheep **Predicted**Human, Mouse, Rat, Pig, Dog, Bovine, Sheep

Host Rabbit
Clonality Polyclonal
Calculated MW 34349

#### **Additional Information**

**Gene ID** 1628

Alias Symbol DABP

Other Names D site-binding protein, Albumin D box-binding protein, Albumin

D-element-binding protein, Tax-responsive enhancer element-binding protein

302, TaxREB302, DBP

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

**Reconstitution & Storage** Add 50 ul of distilled water. Final anti-DBP antibody concentration is 1 mg/ml

in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C.

Avoid repeat freeze-thaw cycles.

**Precautions** DBP antibody - middle region is for research use only and not for use in

diagnostic or therapeutic procedures.

#### **Protein Information**

Name DBP

**Function** This transcriptional activator recognizes and binds to the sequence

5'-RTTAYGTAAY-3' found in the promoter of genes such as albumin, CYP2A4

and CYP2A5. It is not essential for circadian rhythm generation, but

modulates important clock output genes. May be a direct target for regulation by the circadian pacemaker component clock. May affect circadian period and

sleep regulation.

Cellular Location Nucleus.

#### **Tissue Location**

Ubiquitously expressed. Expressed in the suprachiasmatic nuclei (SCN) and in most peripheral tissues, with a strong circadian rhythmicity

### References

Smith, J.S., et al., (2000) Genomics 64 (1), 44-50Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.

## **Images**

