

Bmp7 antibody - N-terminal region

Rabbit Polyclonal Antibody
Catalog # AI10346

Product Information

Application	WB, IHC
Primary Accession	P18075
Other Accession	NM_001191856 , BAA31853
Reactivity	Human, Mouse, Rat, Pig, Dog, Bovine
Predicted	Human, Mouse, Rat, Pig, Chicken, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	49313

Additional Information

Gene ID	655
Alias Symbol	BMP-7, Bmp7
Other Names	Bone morphogenetic protein 7, BMP-7, Osteogenic protein 1, OP-1, Eptotermin alfa, BMP7, OP1
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-Bmp7 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	Bmp7 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	BMP7
Synonyms	OP1
Function	Growth factor of the TGF-beta superfamily that plays important role in various biological processes, including embryogenesis, hematopoiesis, neurogenesis and skeletal morphogenesis (PubMed: 31208997). Initiates the canonical BMP signaling cascade by associating with type I receptor ACVR1 and type II receptor ACVR2A (PubMed: 12667445 , PubMed: 9748228). Once all three components are bound together in a complex at the cell surface, ACVR2A phosphorylates and activates ACVR1. In turn, ACVR1 propagates signal by phosphorylating SMAD1/5/8 that travel to the nucleus and act as

activators and repressors of transcription of target genes (PubMed:[12478285](#)). For specific functions such as growth cone collapse in developing spinal neurons and chemotaxis of monocytes, also uses BMPR2 as type II receptor (PubMed:[31208997](#)). Can also signal through non-canonical pathways such as P38 MAP kinase signaling cascade that promotes brown adipocyte differentiation through activation of target genes, including members of the SOX family of transcription factors (PubMed:[27923061](#)). Promotes the expression of HAMP, this is repressed by its interaction with ERFE (PubMed:[30097509](#)).


Cellular Location

Secreted.


Tissue Location

Expressed in the kidney and bladder. Lower levels seen in the brain

Images




WB Suggested Anti-Bmp7 Antibody Titration: .2-1 ug/ml
ELISA Titer: 1:625
Positive Control: Rat Lung



Human Skin

BMP7



Brown: BMP7
Blue: DAPI