

NMUR2 antibody - N-terminal region

Rabbit Polyclonal Antibody Catalog # AI10816

Product Information

Application WB

Primary Accession <u>Q9GZQ4</u>

Other Accession <u>NM 020167</u>, <u>NP 064552</u>

Reactivity Human
Predicted Human
Host Rabbit
Clonality Polyclonal
Calculated MW 47696

Additional Information

Gene ID 56923

Alias Symbol FM4, FM-4, TGR1, NMU2R, TGR-1, NMU-R2

Other Names Neuromedin-U receptor 2, NMU-R2, G-protein coupled receptor FM-4,

G-protein coupled receptor TGR-1, NMUR2, NMU2R, TGR1

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

Reconstitution & Storage Add 100 ul of distilled water. Final anti-NMUR2 antibody concentration is 1

mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at

20°C. Avoid repeat freeze-thaw cycles.

Precautions NMUR2 antibody - N-terminal region is for research use only and not for use

in diagnostic or therapeutic procedures.

Protein Information

Name NMUR2

Synonyms NMU2R, TGR1

Function Receptor for the neuromedin-U and neuromedin-S neuropeptides.

Cellular Location Cell membrane; Multi-pass membrane protein.


Tissue Location Predominantly expressed in the CNS, particularly in the medulla oblongata,

pontine reticular formation, spinal cord, and thalamus. High level in testis whereas lower levels are present in a variety of peripheral tissues including the gastrointestinal tract, genitourinary tract, liver, pancreas, adrenal gland,

References

Fernandez (2002) Peptides 23 (9), 1617-1623Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.

Images



WB Suggested Anti-NMUR2 Antibody Titration: 2.5 $\mu g/ml$ Positive Control: HepG2 Whole Cell

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.