

SLC22A1 antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI11904

Product Information


Application	WB, IHC
Primary Accession	Q9NQD4
Other Accession	NM_003057 , NP_003048
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Dog, Horse, Bovine
Predicted	Human, Rat, Pig, Dog, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	61 kDa

Additional Information

Alias Symbol	HOCT1, OCT1, oct1_cds
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 100 ul of distilled water. Final anti-SLC22A1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	SLC22A1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Images



Rabbit Anti-SLC22A1 Antibody
Paraffin Embedded Tissue: Human Intestine
Cellular Data: Epithelial cells of intestinal villas
Antibody Concentration: 4.0-8.0 µg/ml
Magnification: 400X

WB Suggested Anti-SLC22A1 Antibody Titration: 2.5µg/ml
Positive Control: Jurkat cell lysate



Host: Rabbit
Target Name:SLC22A1
Sample Tissue:Jurkat
Lane A: Primary Antibody
Lane B: Primary Antibody + Blocking Peptide
Primary Antibody
Concentration:2.5µg/ml
Peptide Concentration: 8.0µg/ml
Lysate Quantity: 25ug/lane Gel
Concentration: 12%

Citations

- [Antioxidant effect of ascorbic acid against cisplatin-induced nephrotoxicity and P-glycoprotein expression in rats](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.