

FICD antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI12719

Product Information

Application	WB, IHC
Primary Accession	Q9BVA6
Other Accession	NM_007076 , NP_009007
Reactivity	Human, Mouse, Rat, Pig, Dog, Guinea Pig, Horse, Bovine
Predicted	Human, Mouse, Rat, Dog, Guinea Pig, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	51778

Additional Information

Gene ID	11153
Alias Symbol	HYPE, MGC5623, UNQ3041, hip13, HIP13
Other Names	Adenosine monophosphate-protein transferase FICD, 2.7.7.n1, AMPylator FICD, FIC domain-containing protein, Huntingtin yeast partner E, Huntingtin-interacting protein 13, HIP-13, Huntingtin-interacting protein E, FICD, HIP13, HYPE
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-FICD antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	FICD antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	FICD (HGNC:18416)
Function	Protein that can both mediate the addition of adenosine 5'- monophosphate (AMP) to specific residues of target proteins (AMPylation), and the removal of the same modification from target proteins (de-AMPylation), depending on the context (By similarity). The side chain of Glu-231 determines which of the two opposing activities (AMPylase or de-AMPylase) will take place (PubMed: 36136088). Acts as a key regulator of the ERN1/IRE1-mediated unfolded protein response (UPR) by mediating AMPylation or de-AMPylation of HSPA5/BiP (PubMed: 25601083 , PubMed: 36136088). In unstressed cells,


acts as an adenylyltransferase by mediating AMPylation of HSPA5/BiP at 'Thr-518', thereby inactivating it (By similarity). In response to endoplasmic reticulum stress, acts as a phosphodiesterase by mediating removal of ATP (de-AMPylation) from HSPA5/BiP at 'Thr-518', leading to restore HSPA5/BiP activity (By similarity). Although it is able to AMPylate RhoA, Rac and Cdc42 Rho GTPases in vitro, Rho GTPases do not constitute physiological substrates (PubMed:[19362538](#), PubMed:[25601083](#)).

Cellular Location	Endoplasmic reticulum membrane; Single-pass type II membrane protein
Tissue Location	Ubiquitous..


References

Clark,H.F.,(2003)GenomeRes.13(10),2265-2270ReconstitutionandStorage:Forshorttermuse,storeat2-8Cupto1 week.Forlongtermstorage,storeat-20Cinsmallaliquotstopreventfreeze-thawcycles.

Images



Human Muscle



WB Suggested Anti-FICD Antibody Titration: 0.2-1 µg/ml
Positive Control: Transfected 293T

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.