

FCRLA antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI13131

Product Information

Application	WB
Primary Accession	Q7L513
Other Accession	NM_032738 , NP_116127
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	38927

Additional Information

Gene ID	84824
Alias Symbol	FCRL, FCRLM1, FCRLX, FCRLb, FCRLc1, FCRLc2, FCRLd, FCRLe, FREB, MGC4595, RP11-474I16.5, FCRX, FCRL1
Other Names	Fc receptor-like A, Fc receptor homolog expressed in B-cells, Fc receptor-like and mucin-like protein 1, Fc receptor-like protein, Fc receptor-related protein X, FcRX, FCRLA, FCRL, FCRL1, FCRLM1, FCRX, FREB
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-FCRLA antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	FCRLA antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Name	FCRLA
Synonyms	FCRL, FCRL1, FCRLM1, FCRX, FREB
Function	May be implicated in B-cell differentiation and lymphomagenesis.
Cellular Location	Cytoplasm. Note=Seems not to be secreted.
Tissue Location	Expressed specifically in primary and secondary lymphoid tissues like lymph node, spleen and tonsil. Specifically expressed in B-cells with a high level in

normal germinal center B- cells, centroblasts and in a subset of diffuse large B-cell lymphomas Highly expressed in bone marrow B-cells and weakly in earlier B lineage cells. Expressed in pre-germinal and germinal center B-cells in secondary lymphoid tissues. Also expressed in melanoma and melanocytes

References

Mechetina L.V.,et al.Eur. J. Immunol. 32:87-96(2002).
Facchetti F.,et al.Proc. Natl. Acad. Sci. U.S.A. 99:3776-3781(2002).
Davis R.S.,et al.Int. Immunol. 14:1075-1083(2002).
Inozume T.,et al.Int. J. Cancer 114:283-290(2005).
Guselnikov S.,et al.Submitted (MAY-2002) to the EMBL/GenBank/DDBJ databases.

Images



WB Suggested Anti-FCRLA Antibody Titration: 0.2-1 µg/ml
ELISA Titer: 1:62500
Positive Control: Hela cell lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.