

MAD2L1BP antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI14660

Product Information

Application	WB
Primary Accession	Q15013
Other Accession	NM_014628 , NP_055443
Reactivity	Human, Pig, Dog, Horse
Predicted	Human, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	31052

Additional Information

Gene ID	9587
Alias Symbol	CMT2, KIAA0110, MGC11282, RP1-261G23.6
Other Names	MAD2L1-binding protein, Caught by MAD2 protein, MAD2L1BP, CMT2, KIAA0110
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-MAD2L1BP antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	MAD2L1BP antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	MAD2L1BP
Synonyms	CMT2, KIAA0110
Function	May function to silence the spindle checkpoint and allow mitosis to proceed through anaphase by binding MAD2L1 after it has become dissociated from the MAD2L1-CDC20 complex.
Cellular Location	Nucleus. Cytoplasm, cytoskeleton, spindle. Note=During early mitosis, unevenly distributed throughout the nucleoplasm. From metaphase to anaphase, concentrated on the spindle

References

Nagase T.,et al.DNA Res. 2:37-43(1995).

Ota T.,et al.Nat. Genet. 36:40-45(2004).

Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.

Mungall A.J.,et al.Nature 425:805-811(2003).

Li W.B.,et al.Submitted (APR-2003) to the EMBL/GenBank/DDBJ databases.

Images



WB Suggested Anti-MAD2L1BP Antibody Titration: 1.0 μ g/ml

Positive Control: MDA-MB-435S Whole CellMAD2L1BP is supported by BioGPS gene expression data to be expressed in MDA-MB435

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.