10320 Camino Santa Fe, Suite G San Diego, CA 92121 Tel: 858.875.1900 Fax: 858.875.1999



Anti-ADGRL3 / LPHN3 Antibody (C-Terminus)

Rabbit Anti Human Polyclonal Antibody Catalog # ALS17503

Product Information

Application IHC-P **Primary Accession** Q9HAR2

Predicted Human, Mouse, Rat, Hamster, Monkey, Chicken, Horse

Host Rabbit
Clonality Polyclonal
Calculated MW 161812
Concentration (mg/ml) 1 mg/ml

Additional Information

Gene ID 23284

Alias Symbol ADGRL3

Other Names ADGRL3, CIRL3, CIRL-3, Lectomedin 3, Lectomedin-3, Lph3, KIAA0768,

Latrophilin 3, Latrophilin homolog 3 (cow), LPHN3, Cl3 latrotoxin receptor,

Latrophilin-3, LEC3

Target/Specificity Human LPHN3. BLAST analysis of the peptide immunogen showed no

homology with other human proteins.

Reconstitution & Storage Immunoaffinity purified

Precautions Anti-ADGRL3 / LPHN3 Antibody (C-Terminus) is for research use only and not

for use in diagnostic or therapeutic procedures.

Protein Information

Name ADGRL3 {ECO:0000303 | PubMed:35418682,

ECO:0000312 | HGNC:HGNC:20974}

Function Orphan adhesion G-protein coupled receptor (aGPCR), which mediates

synapse specificity (PubMed: 35418682). Ligand binding causes a

conformation change that triggers signaling via guanine nucleotide- binding proteins (G proteins) and modulates the activity of downstream effectors (PubMed:35418682). ADGRL3 is coupled with different classes of G alpha proteins, such as G(12)/G(13), G(s), G(i) or G(q), depending on the context (PubMed:35418682). Coupling to G(12)/G(13) G proteins, which mediates the activation Rho small GTPases is the most efficient (PubMed:35418682).

Following G-protein coupled receptor activation, associates with cell adhesion molecules that are expressed at the surface of adjacent cells to direct synapse specificity (PubMed: 26235030). Specifically mediates the establishment of

Schaffer- collateral synapses formed by CA3-region axons on CA1-region pyramidal neurons in the hippocampus (By similarity). Localizes to postsynaptic spines in excitatory synapses in the S.oriens and S.radiatum and interacts with presynaptic cell adhesion molecules FLRT3 and TENM2, promoting synapse formation (By similarity). Plays a role in the development of glutamatergic synapses in the cortex (By similarity). Important in determining the connectivity rates between the principal neurons in the cortex (By similarity).

Cellular Location

Cell membrane; Multi-pass membrane protein. Postsynaptic cell membrane {ECO:0000250|UniProtKB:Q80TS3}; Multi-pass membrane protein. Cell projection, axon {ECO:0000250|UniProtKB:Q80TS3}. Cell junction {ECO:0000250|UniProtKB:Q80TS3}

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.