

ITGB2 Antibody

Mouse Monoclonal Antibody (Mab)
Catalog # AM2138b

Product Information

Application	WB, E
Primary Accession	P05107
Other Accession	NP_000202.2
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Clone Names	534CT6.2.1
Calculated MW	84791

Additional Information

Gene ID	3689
Other Names	Integrin beta-2, Cell surface adhesion glycoproteins LFA-1/CR3/p150, 95 subunit beta, Complement receptor C3 subunit beta, CD18, ITGB2, CD18, MFI7
Target/Specificity	Purified His-tagged ITGB2 protein(Fragment) was used to produced this monoclonal antibody.
Dilution	WB~~1:500~1000 E~~Use at an assay dependent concentration.
Format	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ITGB2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ITGB2 (HGNC:6155)
Synonyms	CD18, MFI7
Function	Integrin ITGAL:ITGB2 is a receptor for ICAM1, ICAM2 and ICAM3 (PubMed: 1676048 , PubMed: 23775590 , PubMed: 38195629). Integrin

ITGAL:ITGB2 is also a receptor for the secreted form of ubiquitin-like protein ISG15; the interaction is mediated by ITGAL (PubMed:[29100055](#)). Integrins ITGAM:ITGB2 and ITGAX:ITGB2 are receptors for the iC3b fragment of the third complement component and for fibrinogen. Integrin ITGAX:ITGB2 recognizes the sequence G-P-R in fibrinogen alpha-chain. Integrin ITGAM:ITGB2 recognizes P1 and P2 peptides of fibrinogen gamma chain. Integrin ITGAM:ITGB2 is also a receptor for factor X. Integrin ITGAD:ITGB2 is a receptor for ICAM3 and VCAM1 (PubMed:[10438935](#), PubMed:[8777714](#), PubMed:[9841932](#)). Contributes to natural killer cell cytotoxicity (PubMed:[15356110](#)). Involved in leukocyte adhesion and transmigration of leukocytes including T-cells and neutrophils (PubMed:[11812992](#), PubMed:[28807980](#)). Triggers neutrophil transmigration during lung injury through PTK2B/PYK2-mediated activation (PubMed:[18587400](#)). Integrin ITGAL:ITGB2 in association with ICAM3, contributes to apoptotic neutrophil phagocytosis by macrophages (PubMed:[23775590](#)). In association with alpha subunit ITGAM/CD11b, required for CD177-PRTN3-mediated activation of TNF primed neutrophils (PubMed:[21193407](#)). Integrins ITGAX:ITGB2 functions as a receptor of the erythrocyte-specific adhesion molecule ICAM4 and mediates erythrophagocytosis (PubMed:[16985175](#)). Integrins ITGAX:ITGB2 functions as a receptor of the neuron-specific adhesion molecule ICAM5 ensuring neuron cell-leukocyte adhesion (PubMed:[10741396](#)). Integrin ITGAL:ITGB2 functions as a receptor of ICAM1 by acting as a platform at the immunological synapse to translate TCR engagement and density of the ITGAL ligand ICAM1 into graded adhesion (PubMed:[38195629](#)). Integrin ITGAM:ITGB2/MAC-1 complex functions as a signaling receptor for the ligand receptor ICAM1, ensuring adhesion between stimulated neutrophils and stimulated endothelial cells (PubMed:[1980124](#)). Integrin ITGAL/ITGB2 that functions as a signaling receptor of ICAM2, ensuring leukocyte cell-cell adhesion on resting cells (PubMed:[1676048](#)).

Cellular Location

Cell membrane; Single-pass type I membrane protein. Membrane raft; Single-pass type I membrane protein

Tissue Location

Leukocytes (PubMed:[23775590](#)). Expressed in neutrophils (at protein level) (PubMed:[21193407](#), PubMed:[28807980](#))


Background

The product of this gene belongs to the integrin beta chain family of proteins. Integrins are integral cell-surface proteins composed of an alpha chain and a beta chain. This gene encodes the integrin beta chain beta 2. A given chain may combine with multiple partners resulting in different integrins. For example, beta 2 combines with the alpha L chain to form the integrin LFA-1, and combines with the alpha M chain to form the integrin Mac-1. Integrins are known to participate in cell adhesion as well as cell-surface mediated signalling. Defects in this gene are the cause of leukocyte adhesion deficiency type I (LAD1). Two transcript variants encoding the same protein have been identified for this gene.

References

- Gjelstrup, L.C., et al. *J. Immunol.* 185(7):4154-4168(2010)
Shimada, M., et al. *Hum. Genet.* 128(4):433-441(2010)
Bailey, S.D., et al. *Diabetes Care* 33(10):2250-2253(2010)
Chen, X., et al. *Proc. Natl. Acad. Sci. U.S.A.* 107(33):14727-14732(2010)
Pliyev, B.K., et al. *Biochem. Biophys. Res. Commun.* 397(2):277-282(2010)

Images



All lanes : Anti-ITGB2 Antibody at 1:2000 dilution + HL-60 cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 90 kDa
Blocking/Dilution buffer: 5% NFDm/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.