

# PIN1 Antibody [Knockdown Validated]

Mouse Monoclonal Antibody (Mab) Catalog # AM2212b

## **Product Information**

**Application** WB, IHC-P, E **Primary Accession** Q13526

**Reactivity** Human, Mouse, Rat, Green Monkey

Host Mouse
Clonality Monoclonal
Isotype IgG1
Clone Names 855CT1.7.5
Calculated MW 18243

# **Additional Information**

Gene ID 5300

Other Names Peptidyl-prolyl cis-trans isomerase NIMA-interacting 1, Peptidyl-prolyl

cis-trans isomerase Pin1, PPIase Pin1, Rotamase Pin1, PIN1

**Target/Specificity** Purified His-tagged PIN1 protein was used to produced this monoclonal

antibody.

**Dilution** WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.

**Format** Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein G column, followed by dialysis

against PBS.

**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** PIN1 Antibody [Knockdown Validated] is for research use only and not for use

in diagnostic or therapeutic procedures.

### **Protein Information**

Name PIN1

**Function** Peptidyl-prolyl cis/trans isomerase (PPIase) that binds to and isomerizes

specific phosphorylated Ser/Thr-Pro (pSer/Thr-Pro) motifs

(PubMed:<u>21497122</u>, PubMed:<u>23623683</u>, PubMed:<u>29686383</u>). By inducing conformational changes in a subset of phosphorylated proteins, acts as a molecular switch in multiple cellular processes (PubMed:<u>21497122</u>, PubMed:<u>22033920</u>, PubMed:<u>23623683</u>). Displays a preference for acidic

residues located N-terminally to the proline bond to be isomerized. Regulates mitosis presumably by interacting with NIMA and attenuating its mitosis-promoting activity. Down-regulates kinase activity of BTK (PubMed:16644721). Can transactivate multiple oncogenes and induce centrosome amplification, chromosome instability and cell transformation. Required for the efficient dephosphorylation and recycling of RAF1 after mitogen activation (PubMed: 15664191). Binds and targets PML and BCL6 for degradation in a phosphorylation-dependent manner (PubMed: 17828269). Acts as a regulator of JNK cascade by binding to phosphorylated FBXW7, disrupting FBXW7 dimerization and promoting FBXW7 autoubiquitination and degradation: degradation of FBXW7 leads to subsequent stabilization of JUN (PubMed:22608923). May facilitate the ubiquitination and proteasomal degradation of RBBP8/CtIP through CUL3/KLHL15 E3 ubiquitin-protein ligase complex, hence favors DNA double-strand repair through error-prone non-homologous end joining (NHEJ) over error-free, RBBP8-mediated homologous recombination (HR) (PubMed:23623683, PubMed:27561354). Upon IL33-induced lung inflammation, catalyzes cis-trans isomerization of phosphorylated IRAK3/IRAK-M, inducing IRAK3 stabilization, nuclear translocation and expression of pro-inflammatory genes in dendritic cells (PubMed: <u>29686383</u>). Catalyzes cis-trans isomerization of phosphorylated phosphoglycerate kinase PGK1 under hypoxic conditions to promote its binding to the TOM complex and targeting to the mitochondrion (PubMed:<u>26942675</u>).

#### **Cellular Location**

Nucleus. Nucleus speckle. Cytoplasm Note=Colocalizes with NEK6 in the nucleus (PubMed:16476580). Mainly localized in the nucleus but phosphorylation at Ser-71 by DAPK1 results in inhibition of its nuclear localization (PubMed:21497122)

#### **Tissue Location**

Expressed in immune cells in the lung (at protein level) (PubMed:29686383). The phosphorylated form at Ser-71 is expressed in normal breast tissue cells but not in breast cancer cells

# **Background**


Essential PPIase that regulates mitosis presumably by interacting with NIMA and attenuating its mitosis-promoting activity. Displays a preference for an acidic residue N-terminal to the isomerized proline bond. Catalyzes pSer/Thr-Pro cis/trans isomerizations. Down-regulates kinase activity of BTK. Can transactivate multiple oncogenes and induce centrosome amplification, chromosome instability and cell transformation. Required for the efficient dephosphorylation and recycling of RAF1 after mitogen activation.


## References

Ebert L., et al. Submitted (MAY-2004) to the EMBL/GenBank/DDBJ databases. Lu K.P., et al. Nature 380:544-547(1996). Kalnine N., et al. Submitted (OCT-2004) to the EMBL/GenBank/DDBJ databases. Ota T., et al. Nat. Genet. 36:40-45(2004). Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.


# **Images**

Immunohistochemical analysis of paraffin-embedded H. breast section using PIN1 Antibody(Cat#AM2212B). AM2212B was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.





All lanes: Anti-PIN1 at 1:2000 dilution Lane 1: K562 whole cell lysate Lane 2: Hela whole cell lysate Lane 3: 293 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 18 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes: Anti-PIN1 Antibody at 1:2000 dilution Lane 1: Hela Lane 2: Hela-Knockdown Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Mouse IgG, (H+L), Peroxidase conjugated (ASP1613) at 1/8000 dilution. Predicted band size: 18 kDa

# **Citations**

- Hyperthermia depletes Oct4 in mouse blastocysts and stem cells
- RACK1 Promotes Self-Renewal and Chemoresistance of Cancer Stem Cells in Human Hepatocellular Carcinoma through Stabilizing Nanog.
- Knockdown of the prolyl isomerase Pin1 inhibits Hep-2 cells growth, migration and invasion by targeting β-catenin signaling pathway.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.