

# MAFK Antibody

Purified Mouse Monoclonal Antibody (Mab)

Catalog # AM8420b

## Product Information

---

<b>Application</b>	WB, FC, E
<b>Primary Accession</b>	<a href="#">O60675</a>
<b>Reactivity</b>	Human, Rat, Mouse
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	IgG1, $\kappa$
<b>Clone Names</b>	1328CT786.105.125
<b>Calculated MW</b>	17523

## Additional Information

---

<b>Gene ID</b>	7975
<b>Other Names</b>	Transcription factor MafK, Erythroid transcription factor NF-E2 p18 subunit, MAFK
<b>Target/Specificity</b>	This MAFK antibody is generated from a mouse immunized with a KLH conjugated synthetic peptide between amino acids from the human region of human MAFK.
<b>Dilution</b>	WB~~1:1000 FC~~1:25 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	MAFK Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	MAFK
<b>Function</b>	Since they lack a putative transactivation domain, the small Mafs behave as transcriptional repressors when they dimerize among themselves (PubMed: <a href="#">9150357</a> ). However, they act as transcriptional activators by dimerizing with other (usually larger) basic-zipper proteins, such as NFE2, NFE2L1/NRF1, NFE2L2/NRF2 and NFE2L3/NRF3, and recruiting them to

specific DNA-binding sites (PubMed:[8932385](#), PubMed:[9150357](#)). Small Maf proteins heterodimerize with Fos and may act as competitive repressors of the NF-E2 transcription factor (PubMed:[9150357](#)).

## Cellular Location

Nucleus.


## Background

Since they lack a putative transactivation domain, the small Mafs behave as transcriptional repressors when they dimerize among themselves. However, they seem to serve as transcriptional activators by dimerizing with other (usually larger) basic-zipper proteins and recruiting them to specific DNA-binding sites. Small Maf proteins heterodimerize with Fos and may act as competitive repressors of the NF-E2 transcription factor.


## References

Toki T.,et al.Oncogene 14:1901-1910(1997).  
Hillier L.W.,et al.Nature 424:157-164(2003).  
Daub H.,et al.Mol. Cell 31:438-448(2008).  
Dephoure N.,et al.Proc. Natl. Acad. Sci. U.S.A. 105:10762-10767(2008).  
Mayya V.,et al.Sci. Signal. 2:RA46-RA46(2009).

## Images



Flow cytometric analysis of HeLa cells using MAFK Antibody(green, Cat#AM8420b) compared to an isotype control of mouse IgG1(blue). AM8420b was diluted at 1:25 dilution. An Alexa Fluor® 488 goat anti-mouse IgG at 1:400 dilution was used as the secondary antibody.



Western blot analysis of lysate from Jurkat cell line, using MAFK Antibody(Cat. #AM8420b). AM8420b was diluted at 1:1000. A goat anti-mouse IgG H&L(HRP) at 1:3000 dilution was used as the secondary antibody. Lysate at 35µg.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.