

RAB3B Antibody

Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM8487b

Product Information

Application	WB, IHC-P, FC, E
Primary Accession	P20337
Reactivity	Human, Rat, Mouse
Host	Mouse
Clonality	monoclonal
Isotype	IgG1,k
Clone Names	1543CT354.60.92
Calculated MW	24758

Additional Information

Gene ID	5865
Other Names	Ras-related protein Rab-3B, RAB3B
Target/Specificity	This RAB3B antibody is generated from a mouse immunized with a recombinant protein of human RAB3B.
Dilution	WB~~1:2000 IHC-P~~1:100~500 FC~~1:25 E~~Use at an assay dependent concentration.
Format	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	RAB3B Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	RAB3B (HGNC:9778)
Function	The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes (PubMed: 35871249). Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different sets of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion (PubMed: 35871249).

Cellular Location

Cell membrane; Lipid-anchor; Cytoplasmic side. Golgi apparatus {ECO:0000250|UniProtKB:Q9CZT8}. Note=Colocalizes with DRC4 in the Golgi apparatus. {ECO:0000250|UniProtKB:Q9CZT8}


Background

Protein transport. Probably involved in vesicular traffic (By similarity).


References

Zahraoui A., et al. *J. Biol. Chem.* 264:12394-12401(1989).
Puhl H.L. III, et al. Submitted (APR-2002) to the EMBL/GenBank/DDBJ databases.
Gregory S.G., et al. *Nature* 441:315-321(2006).
Rigbolt K.T., et al. *Sci. Signal.* 4:RS3-RS3(2011).


Images



Overlay histogram showing HepG2 cells stained with AM8487b (green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AM8487b, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Mouse IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(NA168821) at 1/400 dilution for 40 min at 37°C. Isotype control antibody (blue line) was mouse IgG (1 µg/1x10⁶ cells) used under the same conditions. Acquisition of >10,000 events was performed.



AM8487b staining RAB3B in human pancreas sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hour at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



All lanes : Anti-RAB3B Antibody at 1:2000 dilution Lane 1: mouse brain lysate Lane 2: rat brain lysate Lane 3: SW480 whole cell lysate Lane 4: human brain lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 25 kDa Blocking/Dilution buffer: 5% NFDN/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.