

TPM3 Antibody

Purified Mouse Monoclonal Antibody (Mab)

Catalog # AM8540b

Product Information

Application	WB, E
Primary Accession	P06753
Reactivity	Human, Mouse
Host	Mouse
Clonality	monoclonal
Isotype	IgG1,k
Clone Names	1649CT306.29.13
Calculated MW	32950

Additional Information

Gene ID	7170
Other Names	Tropomyosin alpha-3 chain, Gamma-tropomyosin, Tropomyosin-3, Tropomyosin-5, hTM5, TPM3
Target/Specificity	This TPM3 antibody is generated from a mouse immunized with a recombinant protein of human TPM3.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	TPM3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TPM3
Function	Binds to actin filaments in muscle and non-muscle cells. Plays a central role, in association with the troponin complex, in the calcium dependent regulation of vertebrate striated muscle contraction. Smooth muscle contraction is regulated by interaction with caldesmon. In non-muscle cells is implicated in stabilizing cytoskeleton actin filaments.

Cellular Location

Cytoplasm, cytoskeleton.

Background

Binds to actin filaments in muscle and non-muscle cells. Plays a central role, in association with the troponin complex, in the calcium dependent regulation of vertebrate striated muscle contraction. Smooth muscle contraction is regulated by interaction with caldesmon. In non-muscle cells is implicated in stabilizing cytoskeleton actin filaments.

References

Reinach F.C.,et al.Nature 322:648-650(1986).


McLeod A.R.,et al.Nucleic Acids Res. 14:8413-8426(1986).

Clayton L.,et al.J. Mol. Biol. 201:507-515(1988).

Lin J.J.-C.,et al.Submitted (JUL-2000) to the EMBL/GenBank/DDBJ databases.

Gregory S.G.,et al.Nature 441:315-321(2006).

Images



All lanes : Anti-TPM3 Antibody at 1:2000 dilution Lane 1: HeLa whole cell lysate Lane 2: NIH/3T3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 33 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.