

PFKP Antibody (C-term)

Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM8580b

Product Information

Application	WB, E
Primary Accession	Q01813
Reactivity	Human, Green Monkey, Mouse
Host	Mouse
Clonality	monoclonal
Isotype	IgG1,k
Clone Names	1040CT4.6.4.4
Calculated MW	85596

Additional Information

Gene ID	5214
Other Names	ATP-dependent 6-phosphofructokinase, platelet type {ECO:0000255 HAMAP-Rule:MF_03184}, ATP-PFK {ECO:0000255 HAMAP-Rule:MF_03184}, PFK-P, 2.7.1.11 {ECO:0000255 HAMAP-Rule:MF_03184}, 6-phosphofructokinase type C, Phosphofructo-1-kinase isozyme C, PFK-C, Phosphohexokinase {ECO:0000255 HAMAP-Rule:MF_03184}, PFKP, PFKF
Target/Specificity	This PFKP antibody is generated from a mouse immunized with a recombinant protein between 40-300 amino acids from the C-terminal region of human PFKP.
Dilution	WB~~1:4000 E~~Use at an assay dependent concentration.
Format	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	PFKP Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PFKP (HGNC:8878)
Synonyms	PFKF

Function Catalyzes the phosphorylation of D-fructose 6-phosphate to fructose 1,6-bisphosphate by ATP, the first committing step of glycolysis.

Cellular Location Cytoplasm {ECO:0000255 | HAMAP-Rule:MF_03184}.


Background

Catalyzes the phosphorylation of D-fructose 6-phosphate to fructose 1,6-bisphosphate by ATP, the first committing step of glycolysis.

References

Eto K.,et al.Biochem. Biophys. Res. Commun. 198:990-998(1994).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Deloukas P.,et al.Nature 429:375-381(2004).
Simpson C.J.,et al.Biochem. Biophys. Res. Commun. 180:197-203(1991).
Rush J.,et al.Nat. Biotechnol. 23:94-101(2005).

Images



All lanes : Anti-PFKP Antibody (C-term) at 1:4000 dilution
Lane 1: Hela whole cell lysate Lane 2: Jurkat whole cell lysate Lane 3: COS-7 whole cell lysate Lane 4: mouse heart lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 86 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.