

# PDCD1LG2 Antibody (N-term)

Purified Mouse Monoclonal Antibody (Mab)

Catalog # AM8678b

## Product Information

---

|                          |                        |
|--------------------------|------------------------|
| <b>Application</b>       | WB, IHC-P, E           |
| <b>Primary Accession</b> | <a href="#">Q9BQ51</a> |
| <b>Reactivity</b>        | Human                  |
| <b>Predicted</b>         | Human                  |
| <b>Host</b>              | Mouse                  |
| <b>Clonality</b>         | monoclonal             |
| <b>Isotype</b>           | IgG1, $\kappa$         |
| <b>Clone Names</b>       | 2011CT786.31.38        |
| <b>Calculated MW</b>     | 30957                  |

## Additional Information

---

|                           |  |
|---------------------------|--|
| <b>Gene ID</b>            | 80380  |
| <b>Other Names</b>        | Programmed cell death 1 ligand 2, PD-1 ligand 2, PD-L2, PDCD1 ligand 2, Programmed death ligand 2, Butyrophilin B7-DC, B7-DC, CD273, PDCD1LG2, B7DC, CD273, PDCD1L2, PDL2  |
| <b>Target/Specificity</b> | This PDCD1LG2 antibody is generated from a mouse immunized with a KLH conjugated synthetic peptide between 40-75 amino acids from the N-terminal region of human PDCD1LG2. |
| <b>Dilution</b>           | WB~~1:8000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.  |
| <b>Format</b>             | Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.        |
| <b>Storage</b>            | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.                                    |
| <b>Precautions</b>        | PDCD1LG2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.   |

## Protein Information

---

|                 |   |
|-----------------|---|
| <b>Name</b>     | PDCD1LG2  |
| <b>Synonyms</b> | B7DC, CD273, PDCD1L2, PDL2  |
| <b>Function</b> | Plays a critical role in induction and maintenance of immune tolerance to |

self (PubMed:[11224527](#), PubMed:[12538684](#)). Acts as a ligand for the inhibitory receptor PDCD1/PD-1, inhibiting T-cell proliferation by blocking cell cycle progression and cytokine production (PubMed:[11224527](#)).

### Cellular Location

[Isoform 3]: Secreted [Isoform 1]: Cell membrane; Single-pass type I membrane protein

### Tissue Location

Highly expressed in heart, placenta, pancreas, lung and liver and weakly expressed in spleen, lymph nodes and thymus


## Background

Involved in the costimulatory signal, essential for T- cell proliferation and IFNG production in a PDCD1-independent manner. Interaction with PDCD1 inhibits T-cell proliferation by blocking cell cycle progression and cytokine production (By similarity).


## References

- Tseng S.-Y.,et al.J. Exp. Med. 193:839-846(2001).  
Latchman Y.,et al.Nat. Immunol. 2:261-268(2001).  
He X.-H.,et al.Acta Biochim. Biophys. Sin. 36:284-289(2004).  
Humphray S.J.,et al.Nature 429:369-374(2004).  
Zhang Z.,et al.Protein Sci. 13:2819-2824(2004).

## Images



Immunohistochemical analysis of PDL-2 in untransfected(left) or transfected(right) with 293T cell sections. Cell was fixed with formaldehyde; antigen retrieval was by heat mediation with a EDTA buffer (pH9.0). Samples were incubated with primary antibody (1:25) for 1 hours at room temperature. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



All lanes : Anti-PDCD1LG2 Antibody (N-term) at 1:8000 dilution  
Lane 1: Non-transfected 293T whole cell lysate  
Lane 2: Transfected PD-L2-transfected 293T whole cell lysate  
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 31 kDa  
Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.