

Anti-TRPM8 (Extracellular region) Antibody

Catalog # AN1997

Product Information

Application	WB, ICC
Primary Accession	Q7Z2W7
Host	Rabbit
Clonality	Rabbit Polyclonal
Isotype	IgG
Calculated MW	127685

Additional Information



Gene ID	79054
Other Names	LTrpC6, TRPp8, TRP, TRPM8,
Dilution	WB~~1:1000 ICC~~N/A
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Anti-TRPM8 (Extracellular region) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.
Shipping	Blue Ice

Background

The Transient Receptor Potential Melastatin (TRPM) subfamily of cation-permeable channels is ubiquitous in mammalian tissues. This family includes TRPM1-8. In addition to acting as a calcium-permeant channel, some TRPM family members, TRPM6 and TRPM7, possess serine/threonine kinase activity and autophosphorylation. TRPM8 is thermoactivated at mildly cold temperatures (>25°C), and can also be activated by compounds that cause a cooling sensation, such as menthol and icilin. TRPM8 is expressed in trigeminal and dorsal root ganglia neurons where it confers sensitivity to cold in the somatosensory system. In vascular smooth muscle, TRPM8 may alter blood flow by constricting or enlarging blood vessels. TRPM8 is also expressed in normal prostate epithelial cells, as well as overexpressed in several primary tumors including colon, lung, skin, breast, and prostate cancers.

Images

Western blot image of human TRPM8 in human MDA-MB-231 cells. The blot was probed with rabbit polyclonal anti-TRPM8 (extracellular region) without peptide blocking (lane 1) or with peptide blocking (lane 2).



Immunocytochemical labeling of TRPM8 in paraformaldehyde fixed and NP-40 permeabilized MCF-7 cells. The cells were labeled with rabbit polyclonal anti-TRPM8 (AN1997). The antibody was detected using goat anti-rabbit DyLight® 594.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.