

# IL-10 Antibody

Purified Mouse Monoclonal Antibody  
Catalog # AO1011a

## Product Information

---

<b>Application</b>	WB, E
<b>Primary Accession</b>	<a href="#">P22301</a>
<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Clone Names</b>	3C12C12
<b>Isotype</b>	IgG1
<b>Calculated MW</b>	20517
<b>Description</b>	Interleukine 10 (IL-10) is a cytokine produced primarily by monocytes and to a lesser extent by lymphocytes. This cytokine has pleiotropic effects in immunoregulation and inflammation. It down-regulates the expression of Th1 cytokines, MHC class II Ags, and costimulatory molecules on macrophages. It also enhances B cell survival, proliferation, and antibody production. This cytokine can block NF-kappa B activity, and is involved in the regulation of the JAK-STAT signaling pathway. Knockout studies in mice suggested the function of this cytokine as an essential immunoregulator in the intestinal tract.
<b>Immunogen</b>	Purified recombinant fragment of human IL-10 expressed in E. Coli.
<b>Formulation</b>	Purified antibody in PBS containing 0.03% sodium azide.

## Additional Information

---

<b>Gene ID</b>	3586
<b>Other Names</b>	Interleukin-10, IL-10, Cytokine synthesis inhibitory factor, CSIF, IL10
<b>Dilution</b>	WB~~1/500 - 1/2000 E~~N/A
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	IL-10 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	IL10
<b>Function</b>	Major immune regulatory cytokine that acts on many cells of the immune

system where it has profound anti-inflammatory functions, limiting excessive tissue disruption caused by inflammation. Mechanistically, IL10 binds to its heterotetrameric receptor comprising IL10RA and IL10RB leading to JAK1 and STAT2-mediated phosphorylation of STAT3 (PubMed:[16982608](#)). In turn, STAT3 translocates to the nucleus where it drives expression of anti-inflammatory mediators (PubMed:[18025162](#)). Targets antigen-presenting cells (APCs) such as macrophages and monocytes and inhibits their release of pro-inflammatory cytokines including granulocyte-macrophage colony-stimulating factor /GM-CSF, granulocyte colony-stimulating factor/G-CSF, IL-1 alpha, IL-1 beta, IL-6, IL-8 and TNF (PubMed:[11564774](#), PubMed:[1940799](#), PubMed:[7512027](#)). Also interferes with antigen presentation by reducing the expression of MHC-class II and co-stimulatory molecules, thereby inhibiting their ability to induce T cell activation (PubMed:[8144879](#)). In addition, controls the inflammatory response of macrophages by reprogramming essential metabolic pathways including mTOR signaling (By similarity).

**Cellular Location**

Secreted.

**Tissue Location**

Produced by a variety of cell lines, including T- cells, macrophages, mast cells and other cell types

**References**

---

1. Vieira P, et al. PNAS, 1991.88:1172-1176.

**Images**

---




Figure 1: Western blot analysis using IL10 mouse mAb against IL10 recombinant protein.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.