

EphB4 Antibody

Purified Mouse Monoclonal Antibody

Catalog # AO1184a

Product Information

Application	WB, E
Primary Accession	P54760
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Clone Names	7H4A6
Isotype	IgG1
Calculated MW	108270
Description	EphB4: EPH receptor B4. Ephrin receptors and their ligands, the ephrins, mediate numerous developmental processes, particularly in the nervous system. Based on their structures and sequence relationships, ephrins are divided into the ephrin-A (EFNA) class, which are anchored to the membrane by a glycosylphosphatidylinositol linkage, and the ephrin-B (EFNB) class, which are transmembrane proteins. The Eph family of receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. Ephrin receptors make up the largest subgroup of the receptor tyrosine kinase (RTK) family. The protein encoded by this gene binds to ephrin-B2 and plays an essential role in vascular development.
Immunogen	Purified recombinant fragment of EphB4 (aa562-612)expressed in E. Coli.
Formulation	Ascitic fluid containing 0.03% sodium azide.

Additional Information

Gene ID	2050
Other Names	Ephrin type-B receptor 4, 2.7.10.1, Hepatoma transmembrane kinase, Tyrosine-protein kinase TYRO11, EPHB4, HTK, MYK1, TYRO11
Dilution	WB~~1/500 - 1/2000 E~~N/A
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	EphB4 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	EPHB4
Synonyms	HTK, MYK1, TYRO11
Function	Receptor tyrosine kinase which binds promiscuously transmembrane ephrin-B family ligands residing on adjacent cells, leading to contact-dependent bidirectional signaling into neighboring cells. The signaling pathway downstream of the receptor is referred to as forward signaling while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling. Together with its cognate ligand/functional ligand EFNB2 it is involved in the regulation of cell adhesion and migration, and plays a central role in heart morphogenesis, angiogenesis and blood vessel remodeling and permeability. EPHB4-mediated forward signaling controls cellular repulsion and segregation from EFNB2-expressing cells.
Cellular Location	Cell membrane; Single-pass type I membrane protein
Tissue Location	Abundantly expressed in placenta but also detected in kidney, liver, lung, pancreas, skeletal muscle and heart. Expressed in primitive and myeloid, but not lymphoid, hematopoietic cells. Also observed in cell lines derived from liver, breast, colon, lung, melanocyte and cervix.

References

1. Biochem Biophys Res Commun. 2004 Aug 27;321(3):681-7. 2. Cancer Res. 2005 Jun 1;65(11):4623-32. 3. Development. 2005 Sep;132(18):4097-106.

Images




Figure 1: Western blot analysis using EphB4 mouse mAb against Jurkat (1) and HEK293 (2) cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.