

# KID Antibody

Purified Mouse Monoclonal Antibody

Catalog # AO1625a

## Product Information

---

<b>Application</b>	WB, IHC, E
<b>Primary Accession</b>	<a href="#">Q14807</a>
<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Clone Names</b>	5F3
<b>Isotype</b>	IgG1
<b>Calculated MW</b>	73262
<b>Description</b>	The protein encoded by this gene is a member of kinesin-like protein family. This family of proteins are microtubule-dependent molecular motors that transport organelles within cells and move chromosomes during cell division. The C-terminal half of this protein has been shown to bind DNA. Studies with the Xenopus homolog suggests its essential role in metaphase chromosome alignment and maintenance.
<b>Immunogen</b>	Purified recombinant fragment of human KID expressed in E. Coli.
<b>Formulation</b>	Ascitic fluid containing 0.03% sodium azide.

## Additional Information

---

<b>Gene ID</b>	3835
<b>Other Names</b>	Kinesin-like protein KIF22, Kinesin-like DNA-binding protein, Kinesin-like protein 4, KIF22, KID, KNSL4
<b>Dilution</b>	WB~~1/500 - 1/2000 IHC~~1/200 - 1/1000 E~~1/10000
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	KID Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	KIF22
<b>Synonyms</b>	KID, KNSL4

<b>Function</b>	Kinesin family member that is involved in spindle formation and the movements of chromosomes during mitosis and meiosis. Binds to microtubules and to DNA (By similarity). Plays a role in congression of laterally attached chromosomes in NDC80-depleted cells (PubMed: <a href="#">25743205</a> ).
<b>Cellular Location</b>	Nucleus. Cytoplasm, cytoskeleton
<b>Tissue Location</b>	Expressed in bone, cartilage, joint capsule, ligament, skin, and primary cultured chondrocytes

## References

1. Cell. 2008 Mar 7;132(5):771-82. 2. Retrovirology. 2009 May 19;6:47.

## Images




Figure 1: Western blot analysis using KID mAb against human KID (AA: 225-419) recombinant protein. (Expected MW is 47 kDa)




Figure 2: Western blot analysis using KID mouse mAb against MCF-7 (1) and HeLa (2) cell lysate.





Figure 3: Immunohistochemical analysis of paraffin-embedded cervical cancer tissues using KID mouse mAb with DAB staining.

Figure 4: Immunohistochemical analysis of paraffin-embedded rectum cancer tissues using KID mouse mAb with DAB staining.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.