

FZD6 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP11009c

Product Information

| | |
|--------------------------|---|
| Application | WB, E |
| Primary Accession | O60353 |
| Other Accession | Q61089 , NP_001158087.1 , NP_003497.2 |
| Reactivity | Human |
| Predicted | Mouse |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | Rabbit IgG |
| Clone Names | RB25725 |
| Calculated MW | 79292 |
| Antigen Region | 493-520 |

Additional Information

| | |
|---------------------------|--|
| Gene ID | 8323 |
| Other Names | Frizzled-6, Fz-6, hFz6, FZD6 |
| Target/Specificity | This FZD6 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 493-520 amino acids from the Central region of human FZD6. |
| Dilution | WB~~1:1000 E~~Use at an assay dependent concentration. |
| Format | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| Storage | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| Precautions | FZD6 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures. |

Protein Information

| | |
|-----------------|--|
| Name | FZD6 |
| Function | Receptor for Wnt proteins. Most of frizzled receptors are coupled to the beta-catenin canonical signaling pathway, which leads to the activation of disheveled proteins, inhibition of GSK-3 kinase, nuclear accumulation of |

beta-catenin and activation of Wnt target genes. A second signaling pathway involving PKC and calcium fluxes has been seen for some family members, but it is not yet clear if it represents a distinct pathway or if it can be integrated in the canonical pathway, as PKC seems to be required for Wnt-mediated inactivation of GSK-3 kinase. Both pathways seem to involve interactions with G-proteins. May be involved in transduction and intercellular transmission of polarity information during tissue morphogenesis and/or in differentiated tissues. Together with FZD3, is involved in the neural tube closure and plays a role in the regulation of the establishment of planar cell polarity (PCP), particularly in the orientation of asymmetric bundles of stereocilia on the apical faces of a subset of auditory and vestibular sensory cells located in the inner ear (By similarity).

Cellular Location

Membrane {ECO:0000250|UniProtKB:Q61089}; Multi-pass membrane protein. Cell membrane {ECO:0000250|UniProtKB:Q61089}; Multi-pass membrane protein. Cell surface {ECO:0000250|UniProtKB:Q61089}. Apical cell membrane; Multi-pass membrane protein. Cytoplasmic vesicle membrane {ECO:0000250|UniProtKB:Q61089}; Multi-pass membrane protein. Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:Q61089}; Multi-pass membrane protein. Note=Colocalizes with FZD3 at the apical face of cells (By similarity). Localizes to the endoplasmic reticulum membrane in the presence of LMBR1L (By similarity). {ECO:0000250|UniProtKB:Q61089}

Tissue Location

Detected in adult heart, brain, placenta, lung, liver, skeletal muscle, kidney, pancreas, thymus, prostate, testis, ovary, small intestine and colon. In the fetus, expressed in brain, lung, liver and kidney

Background


This gene represents a member of the 'frizzled' gene family, which encode 7-transmembrane domain proteins that are receptors for Wnt signaling proteins. The protein encoded by this family member contains a signal peptide, a cysteine-rich domain in the N-terminal extracellular region, and seven transmembrane domains, but unlike other family members, this protein does not contain a C-terminal PDZ domain-binding motif. This protein functions as a negative regulator of the canonical Wnt/beta-catenin signaling cascade, thereby inhibiting the processes that trigger oncogenic transformation, cell proliferation, and inhibition of apoptosis. Alternative splicing results in multiple transcript variants.

References

- Yerges, L.M., et al. J. Bone Miner. Res. 24(12):2039-2049(2009)
Kim, J.G., et al. J. Korean Med. Sci. 24(3):443-447(2009)
Miyakoshi, T., et al. Endocr. Pathol. 19(4):261-273(2008)
Sirchia, R., et al. Biol. Chem. 388(5):457-465(2007)
Lyons, J.P., et al. Exp. Cell Res. 298(2):369-387(2004)

Images

FZD6 Antibody (Center) (Cat. #AP11009c) western blot analysis in A549 cell line lysates (35ug/lane). This demonstrates the FZD6 antibody detected the FZD6 protein (arrow).



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.