

JAK2 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1125b

Product Information

Application	WB, IF, FC, E
Primary Accession	O60674
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	130674
Antigen Region	1101-1132

Additional Information

Gene ID	3717
Other Names	Tyrosine-protein kinase JAK2, Janus kinase 2, JAK-2, JAK2
Target/Specificity	This JAK2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1101-1132 amino acids from the C-terminal region of human JAK2.
Dilution	WB~~1:1000 IF~~1:10~50 FC~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	JAK2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	JAK2 (HGNC:6192)
Function	Non-receptor tyrosine kinase involved in various processes such as cell growth, development, differentiation or histone modifications. Mediates essential signaling events in both innate and adaptive immunity. In the cytoplasm, plays a pivotal role in signal transduction via its association with type I receptors such as growth hormone (GHR), prolactin (PRLR), leptin

(LEPR), erythropoietin (EPOR), thrombopoietin receptor (MPL/TPOR); or type II receptors including IFN- alpha, IFN-beta, IFN-gamma and multiple interleukins (PubMed:[15690087](#), PubMed:[7615558](#), PubMed:[9657743](#), PubMed:[15899890](#)). Following ligand- binding to cell surface receptors, phosphorylates specific tyrosine residues on the cytoplasmic tails of the receptor, creating docking sites for STATs proteins (PubMed:[15690087](#), PubMed:[9618263](#)). Subsequently, phosphorylates the STATs proteins once they are recruited to the receptor. Phosphorylated STATs then form homodimer or heterodimers and translocate to the nucleus to activate gene transcription. For example, cell stimulation with erythropoietin (EPO) during erythropoiesis leads to JAK2 autophosphorylation, activation, and its association with erythropoietin receptor (EPOR) that becomes phosphorylated in its cytoplasmic domain (PubMed:[9657743](#)). Then, STAT5 (STAT5A or STAT5B) is recruited, phosphorylated and activated by JAK2. Once activated, dimerized STAT5 translocates into the nucleus and promotes the transcription of several essential genes involved in the modulation of erythropoiesis. Part of a signaling cascade that is activated by increased cellular retinol and that leads to the activation of STAT5 (STAT5A or STAT5B) (PubMed:[21368206](#)). In addition, JAK2 mediates angiotensin-2-induced ARHGEF1 phosphorylation (PubMed:[20098430](#)). Plays a role in cell cycle by phosphorylating CDKN1B (PubMed:[21423214](#)). Cooperates with TEC through reciprocal phosphorylation to mediate cytokine-driven activation of FOS transcription. In the nucleus, plays a key role in chromatin by specifically mediating phosphorylation of 'Tyr-41' of histone H3 (H3Y41ph), a specific tag that promotes exclusion of CBX5 (HP1 alpha) from chromatin (PubMed:[19783980](#)). Up-regulates the potassium voltage- gated channel activity of KCNA3 (PubMed:[25644777](#)).

Cellular Location	Endomembrane system; Peripheral membrane protein. Cytoplasm. Nucleus
Tissue Location	Ubiquitously expressed throughout most tissues.

Background



This gene product is a protein tyrosine kinase involved in a specific subset of cytokine receptor signaling pathways. It has been found to be constitutively associated with the prolactin receptor and is required for responses to gamma interferon. Mice that do not express an active protein for this gene exhibit embryonic lethality associated with the absence of definitive erythropoiesis.

References


- Joos, S., et al., *Int. J. Cancer* 103(4):489-495 (2003).
Leung, K.C., et al., *Proc. Natl. Acad. Sci. U.S.A.* 100(3):1016-1021 (2003).
Saharinen, P., et al., *J. Biol. Chem.* 277(49):47954-47963 (2002).
Giordanetto, F., et al., *Protein Eng.* 15(9):727-737 (2002).
Deo, D.D., et al., *J. Biol. Chem.* 277(24):21237-21245 (2002).

Images

The anti-JAK2 Pab (Cat. #AP1125b) is used in Western blot to detect JAK2 in mouse thymus tissue lysate.



Confocal immunofluorescent analysis of JAK2 Antibody (C-term) (Cat#AP1125b) with Hela cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).



JAK2 Antibody (C-term) (Cat. #AP1125b) flow cytometric analysis of K562 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Citations

- [Evodiamine Induces Apoptosis and Inhibits Migration of HCT-116 Human Colorectal Cancer Cells.](#)
- [Roflumilast reverses polymicrobial sepsis-induced liver damage by inhibiting inflammation in mice.](#)
- [Prolactin enhances interferon-gamma-induced production of CXC ligand 9 \(CXCL9\), CXCL10, and CXCL11 in human keratinocytes.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.