

SUMO2/3 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP1224a

Product Information

Application	IHC-P, WB, IF, E
Primary Accession	P55854
Other Accession	Q7SZ22 , Q5XIF4 , Q9Z172 , Q6DI05 , Q17QV3 , P61959 , P61958 , P61957 , Q2PFW2 , P61956 , Q6DHL4 , Q6LDZ8 , Q5ZJM9 , P61955 , Q6NV25 , Q6GPW2 , Q7ZTK7
Reactivity	Human, Rat, Mouse
Predicted	Xenopus, Zebrafish, Bovine, Chicken, Hamster, Monkey, Pig, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	11637
Antigen Region	49-81

Additional Information

Gene ID	6612
Other Names	Small ubiquitin-related modifier 3, SUMO-3, SMT3 homolog 1 {ECO:0000312 HGNC:HGNC:11124}, SUMO-2, Ubiquitin-like protein SMT3A, Smt3A, SUMO3 (HGNC:11124)
Target/Specificity	This SUMO2/3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 49-81 amino acids from the C-terminal region of human SUMO2/3.
Dilution	IHC-P~~1:100~500 WB~~1:1000 IF~~1:100 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	SUMO2/3 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	SUMO3 (HGNC:11124)
-------------	--------------------------------------

Function	Ubiquitin-like protein which can be covalently attached to target lysines either as a monomer or as a lysine-linked polymer. Does not seem to be involved in protein degradation and may function as an antagonist of ubiquitin in the degradation process. Plays a role in a number of cellular processes such as nuclear transport, DNA replication and repair, mitosis and signal transduction. Covalent attachment to its substrates requires prior activation by the E1 complex SAE1-SAE2 and linkage to the E2 enzyme UBE2I, and can be promoted by an E3 ligase such as PIAS1-4, RANBP2 or CBX4 (PubMed: 11451954 , PubMed: 18538659 , PubMed: 21965678). Plays a role in the regulation of sumoylation status of SETX (PubMed: 24105744).
Cellular Location	Cytoplasm. Nucleus. Nucleus, PML body
Tissue Location	Expressed predominantly in liver.


Background

SUMO2 and SUMO3 are members of the SUMO (small ubiquitin-like modifier) protein family. This protein family functions in a manner similar to ubiquitin in that it is bound to target proteins as part of a post-translational modification system. However, unlike ubiquitin which targets proteins for degradation, this protein is involved in a variety of cellular processes, such as nuclear transport, transcriptional regulation, apoptosis, and protein stability. In vertebrates, three members of the SUMO family have been described, SUMO 1 and the functionally distinct homologues SUMO 2 and SUMO 3. SUMO modification sites present in the N terminal regions of SUMO 2 and SUMO 3 are utilized by SAE1/SAE2 (SUMO E1) and Ubc9 (SUMO E2) to form polymeric chains of SUMO 2 and SUMO 3 on protein substrates, a property not shared by SUMO 1.

References



Strausberg, R.L., et al., Proc. Natl. Acad. Sci. U.S.A. 99(26):16899-16903 (2002).
Lapenta, V., et al., Genomics 40(2):362-366 (1997).

Images



All lanes : Anti-SUMO2/3 Antibody (C-term) at 1:2000 dilution
Lane 1: SH-SY5Y whole cell lysate
Lane 2: HeLa whole cell lysate
Lane 3: Jurkat whole cell lysate
Lane 4: HL-60 whole cell lysate
Lane 5: Mouse cerebellum tissue lysate
Lane 6: Mouse liver tissue lysate
Lane 7: Rat liver tissue lysate
Lysates/proteins at 20 µg per lane.
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 12 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.

Fluorescent image of SH-SY5Y cells stained with SUMO2/3 Antibody (C-term) (Cat#AP1224a). AP1224a was diluted at 1:100 dilution. An Alexa Fluor 488-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody (green). Cytoplasmic actin was counterstained with Alexa Fluor® 555 conjugated with Phalloidin (red).



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

Citations

- [TRIM11 Prevents and Reverses Protein Aggregation and Rescues a Mouse Model of Parkinson's Disease](#)
- [The SUMOylation landscape of renal cortical collecting duct cells](#)
- [HSP70-Hrd1 axis precludes the oncorepressor potential of N-terminal misfolded Blimp-1s in lymphoma cells](#)
- [TRIM3 Promotes APL Progression through Stabilization of the Oncoprotein PML-RAR \$\alpha\$ and Inhibition of p53-Mediated Senescence](#)
- [Adenovirus E4-ORF3 Targets PIAS3 and Together with E1B-55K Remodels SUMO Interactions in the Nucleus and at Virus Genome Replication Domains](#)
- [Signaling via the IL-20 receptor inhibits cutaneous production of IL-1 \$\beta\$ and IL-17A to promote infection with methicillin-resistant *Staphylococcus aureus*](#)
- [PKC \$\zeta\$ mediates disturbed flow-induced endothelial apoptosis via p53 SUMOylation](#)
- [Lysine deacetylation in ischaemic preconditioning: the role of SIRT1](#)
- [Keratin hypersumoylation alters filament dynamics and is a marker for human liver disease and keratin mutation](#)
- [Neuroprotection resulting from insufficiency of RANBP2 is associated with the modulation of protein and lipid homeostasis of functionally diverse but linked pathways in response to oxidative stress](#)
- [Spatial interplay between PIASy and FIP200 in the regulation of signal transduction and transcriptional activity](#)
- [SUMO modification of the Ets-related transcription factor ERM inhibits its transcriptional activity](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.