

# SUMO4 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)  
Catalog # AP1263b

## Product Information

---

<b>Application</b>	IHC-P, WB, E
<b>Primary Accession</b>	<a href="#">Q6EEV6</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB7259

## Additional Information

---

<b>Other Names</b>	Small ubiquitin-related modifier 4, SUMO-4, Small ubiquitin-like protein 4, SUMO4, SMT3H4
<b>Target/Specificity</b>	This SUMO4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide selected from the center region of human SUMO4.
<b>Dilution</b>	IHC-P~~1:100~500 WB~~1:1000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	SUMO4 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

### Background

---


SUMO4 is a member of the SUMO gene family. This family of small ubiquitin-related modifiers covalently modify target lysines in proteins and control the target proteins' subcellular localization, stability, or activity. Upon oxidative stress, SUMO4 conjugates to various anti-oxidant enzymes, chaperones, and stress defense proteins. This protein may also conjugate to NFKBIA, TFAP2A and FOS, negatively regulating their transcriptional activity, and to NR3C1, positively regulating its transcriptional activity. Covalent attachment to SUMO4 substrates requires prior activation by the E1 complex SAE1-SAE2 and linkage to the E2 enzyme UBE2I. In contrast to SUMO1, SUMO2 and SUMO3, SUMO4 seems to be insensitive to sentrin-specific

proteases due to the presence of Pro-90. This may impair processing to mature form and conjugation to substrates. SUMO4 is located in the cytoplasm and specifically modifies IKBA, leading to negative regulation of NF-kappa-B-dependent transcription of the IL12B gene. The M55V substitution has been associated with type I diabetes.


## References

- Park, Y., et al. *Nat. Genet.* 37 (2), 112 (2005)  
Guo, D., et al. *Nat. Genet.* 36 (8), 837-841 (2004)  
Bohren, K.M., et al. *J. Biol. Chem.* 279 (26), 27233-27238 (2004)  
Yang, S.H., et al., *Mol. Cell* 13(4):611-617 (2004).  
Bailey, D., et al., *J. Biol. Chem.* 279(1):692-703 (2004).  
Ling, Y., et al., *Nucleic Acids Res.* 32(2):598-610 (2004).  
Pountney, D.L., et al., *Exp. Neurol.* 184(1):436-446 (2003).  
Ohshima, T., et al., *J. Biol. Chem.* 278(51):50833-50842 (2003).


## Images



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.



Western blot analysis of SUMO4 Antibody (Center) (Cat. #AP1263b) in HepG2 cell line lysate. SUMO4 (arrow) was detected using the purified Pab.



Western blot analysis of SUMO4 (arrow) using rabbit polyclonal SUMO4 Antibody (Center) (Cat. #AP1263b). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the SUMO4 gene.

## Citations

---

- [Cooperative control of striated muscle mass and metabolism by MuRF1 and MuRF2.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.