

# **ELAVL2 Antibody (Center)**

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP12937c

#### **Product Information**

**Application** IHC-P, WB, FC, E

Primary Accession <u>Q12926</u>

Other Accession 075ZT7, 009032, 061701, P26378, 091584, 060900, 014576, 091903,

Q8CH84, Q60899, NP\_001164666.1, NP\_001164668.1

**Reactivity** Human, Mouse **Predicted** Mouse, Rat, Xenopus

HostRabbitClonalityPolyclonalIsotypeRabbit IgGClone NamesRB28546Calculated MW39504Antigen Region156-184

### **Additional Information**

**Gene ID** 1993

Other Names ELAV-like protein 2, ELAV-like neuronal protein 1, Hu-antigen B, HuB, Nervous

system-specific RNA-binding protein Hel-N1, ELAVL2, HUB

Target/Specificity This ELAVL2 antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 156-184 amino acids from the Central

region of human ELAVL2.

Dilution IHC-P~~1:100~500 WB~~1:1000 FC~~1:10~50 E~~Use at an assay dependent

concentration.

**Format** Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** ELAVL2 Antibody (Center) is for research use only and not for use in

diagnostic or therapeutic procedures.

#### **Protein Information**

Name ELAVL2

HUB **Synonyms** 

**Function** RNA-binding protein that binds to the 3' untranslated region (3'UTR) of

target mRNAs (By similarity). Seems to recognize a GAAA motif (By similarity).

Can bind to its own 3'UTR, the FOS 3'UTR and the ID 3'UTR (By similarity).

Brain; neural-specific. **Tissue Location** 


## **Background**

The protein encoded by this gene is a neural-specific RNA-binding protein that is known to bind to several 3' UTRs, including its own and also that of FOS and ID. The encoded protein may recognize a GAAA motif in the RNA. Three transcript variants encoding two different isoforms have been found for this gene.


## References

Marroni, F., et al. Circ Cardiovasc Genet 2(4):322-328(2009) Lowe, J.K., et al. PLoS Genet. 5 (2), E1000365 (2009): D'Alessandro, V., et al. Cell. Oncol. 30(4):291-297(2008) Jonson, L., et al. Mol. Cell Proteomics 6(5):798-811(2007) Yano, M., et al. J. Biol. Chem. 280(13):12690-12699(2005)

## **Images**





Immunohistochemical analysis of AP12937C on paraffin-embedded Human cerebellum tissue. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:100) for 1 hour at room temperature. Undiluted CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



Immunohistochemical analysis of AP12937C on paraffin-embedded Human brain tissue. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:100) for 1 hour at room temperature. Undiluted CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.

Anti-ELAVL2 Antibody (Center) at 1:2000 dilution + Mouse brain lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 40 kDa Blocking/Dilution buffer: 5% NFDM/TBST.





ELAVL2 Antibody (Center) (Cat. #AP12937c) flow cytometric analysis of K562 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.