

NDST1 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP13224b

Product Information

Application WB, IHC-P, E **Primary Accession** P52848

Other Accession <u>Q02353, Q3UHN9, NP 001534.1</u>

Reactivity Human **Predicted** Mouse, Rat Host Rabbit Clonality Polyclonal Isotype Rabbit IgG **Clone Names** RB33030 100868 **Calculated MW** 820-849 **Antigen Region**

Additional Information

Gene ID 3340

Other Names Bifunctional heparan sulfate N-deacetylase/N-sulfotransferase 1,

Glucosaminyl N-deacetylase/N-sulfotransferase 1, NDST-1, N-heparan sulfate

sulfotransferase 1, N-HSST 1, [Heparan sulfate]-glucosamine

N-sulfotransferase 1, HSNST 1, Heparan sulfate N-deacetylase 1, 3---, Heparan

sulfate N-sulfotransferase 1, 282-, NDST1, HSST, HSST1

Target/Specificity This NDST1 antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 820-849 amino acids from the

C-terminal region of human NDST1.

Dilution WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.

Format Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions NDST1 Antibody (C-term) is for research use only and not for use in diagnostic

or therapeutic procedures.

Protein Information

Name NDST1 (HGNC:7680)

Synonyms HSST, HSST1

Function [Isoform 1]: Essential bifunctional enzyme that catalyzes both the

N-deacetylation and the N-sulfation of glucosamine (GlcNAc) of the glycosaminoglycan in heparan sulfate (PubMed:35137078, PubMed:9230113, PubMed:9744796). Modifies the GlcNAc-GlcA disaccharide repeating sugar backbone to make N-sulfated heparosan, a prerequisite substrate for later modifications in heparin biosynthesis (PubMed:9230113). Plays a role in determining the extent and pattern of sulfation of heparan sulfate. Participates in biosynthesis of heparan sulfate that can ultimately serve as L-selectin ligands, thereby playing a role in inflammatory response (By similarity). Required for the exosomal release of SDCBP, CD63 and syndecan

(PubMed:22660413).

Cellular Location [Isoform 1]: Golgi apparatus, trans-Golgi network membrane; Single-pass type

II membrane protein. Golgi apparatus, cis-Golgi network membrane;

Single-pass type II membrane protein

Tissue Location Widely expressed. Expression is most abundant in heart, liver and pancreas.


Background

NDST1 is an essential bifunctional enzyme that catalyzes both the N-deacetylation and the N-sulfation of glucosamine (GlcNAc) of the glycosaminoglycan in heparan sulfate. Modifies the GlcNAc-GlcA dissacharide repeating sugar backbone to make N-sulfated heparosan, a prerequisite substrate for later modifications in heparin biosynthesis. Plays a role in determining the extent and pattern of sulfation of heparan sulfate. Compared to other NDST enzymes, its presence is absolutely required. Participates in biosynthesis of heparan sulfate that can ultimately serve as L-selectin ligands, thereby playing a role in inflammatory response.

References


Yoshida, T., et al. Int. J. Mol. Med. 25(4):649-656(2010) Oguri, M., et al. Am. J. Hypertens. 23(1):70-77(2010) Duelli, A., et al. J. Immunol. 183(11):7073-7083(2009) Zuberi, R.I., et al. J. Immunol. 183(6):3971-3979(2009) Drenos, F., et al. Hum. Mol. Genet. 18(12):2305-2316(2009)

Images



NDST1 Antibody (C-term) (Cat. #AP13224b) western blot analysis in CEM cell line lysates (35ug/lane). This demonstrates the NDST1 antibody detected the NDST1 protein (arrow).

NDST1 Antibody (C-term) (Cat. #AP13224b)immunohistochemistry analysis in formalin fixed and paraffin embedded human liver tissue followed



by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of NDST1 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.