

GFRA4 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP13249c

Product Information

Application WB, E **Primary Accession** Q9GZZ7

Other Accession NP 665705.1, NP 071422.1

Reactivity Human
Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Clone Names RB33123
Calculated MW 31670
Antigen Region 156-184

Additional Information

Gene ID 64096

Other Names GDNF family receptor alpha-4, GDNF receptor alpha-4, GDNFR-alpha-4,

GFR-alpha-4, Persephin receptor, GFRA4

Target/Specificity This GFRA4 antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 156-184 amino acids from the Central

region of human GFRA4.

Dilution WB~~1:1000 E~~Use at an assay dependent concentration.

Format Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions GFRA4 Antibody (Center) is for research use only and not for use in diagnostic

or therapeutic procedures.

Protein Information

Name GFRA4

Function Receptor for persephin (PSPN), a growth factor that exhibits neurotrophic

activity on mesencephalic dopaminergic and motor neurons

(PubMed: 11116144). Acts by binding to its coreceptor, GFRA4, leading to

autophosphorylation and activation of the RET receptor (PubMed:<u>11116144</u>). May be important in C-cell development and, in the postnatal development of

the adrenal medulla (PubMed:11116144).

Cellular Location [Isoform GFRalpha4a]: Cell membrane; Lipid-anchor, GPI-anchor [Isoform

GFRalpha4c]: Secreted

Tissue Location Predominantly expressed in the adult thyroid gland. Low levels also found in

fetal adrenal and thyroid glands


Background

The protein encoded by this gene is a member of the GDNF receptor family. It is a glycosylphosphatidylinositol(GPI)-linked cell surface receptor for persephin, and mediates activation of the RET tyrosine kinase receptor. This gene is a candidate gene for RET-associated diseases. Alternatively spliced transcript variants encoding different isoforms have been described for this gene.

References

Yang, J., et al. FEBS Lett. 569 (1-3), 267-271 (2004):
Borrego, S., et al. J. Med. Genet. 40 (3), E18 (2003):
Deloukas, P., et al. Nature 414(6866):865-871(2001)
Zhou, B., et al. Pediatr. Neurol. 25(2):156-161(2001)
Lindahl, M., et al. J. Biol. Chem. 276(12):9344-9351(2001)

Images



GFRA4 Antibody (Center) (Cat. #AP13249c) western blot analysis in HL-60,ZR-75-1,MDA-MB453 cell line lysates (35ug/lane). This demonstrates the GFRA4 antibody detected the GFRA4 protein (arrow).

Citations

• <u>Downregulation of microRNA-483-5p Promotes Cell Proliferation and Invasion by Targeting GFRA4 in Hirschsprung's Disease.</u>

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.