

TARDBP Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP13763a

Product Information

Application	IHC-P, IF, WB, E
Primary Accession	Q13148
Other Accession	Q921F2 , Q5ZLN5 , NP_031401.1
Reactivity	Human, Rat, Mouse
Predicted	Chicken, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	44740
Antigen Region	1-30

Additional Information

Gene ID	23435
Other Names	TAR DNA-binding protein 43, TDP-43, TARDBP, TDP43
Target/Specificity	This TARDBP antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human TARDBP.
Dilution	IHC-P~~1:100~500 IF~~1:10~50 WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	TARDBP Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TARDBP {ECO:0000303 PubMed:18396105, ECO:0000312 HGNC:HGNC:11571}
Function	RNA-binding protein that is involved in various steps of RNA biogenesis and processing (PubMed: 23519609). Preferentially binds, via its two RNA

recognition motifs RRM1 and RRM2, to GU-repeats on RNA molecules predominantly localized within long introns and in the 3'UTR of mRNAs (PubMed:[23519609](#), PubMed:[24240615](#), PubMed:[24464995](#)). In turn, regulates the splicing of many non-coding and protein-coding RNAs including proteins involved in neuronal survival, as well as mRNAs that encode proteins relevant for neurodegenerative diseases (PubMed:[21358640](#), PubMed:[29438978](#)). Plays a role in maintaining mitochondrial homeostasis by regulating the processing of mitochondrial transcripts (PubMed:[28794432](#)). Also regulates mRNA stability by recruiting CNOT7/CAF1 deadenylase on mRNA 3'UTR leading to poly(A) tail deadenylation and thus shortening (PubMed:[30520513](#)). In response to oxidative insult, associates with stalled ribosomes localized to stress granules (SGs) and contributes to cell survival (PubMed:[19765185](#), PubMed:[23398327](#)). Also participates in the normal skeletal muscle formation and regeneration, forming cytoplasmic myo-granules and binding mRNAs that encode sarcomeric proteins (PubMed:[30464263](#)). Plays a role in the maintenance of the circadian clock periodicity via stabilization of the CRY1 and CRY2 proteins in a FBXL3-dependent manner (PubMed:[27123980](#)). Negatively regulates the expression of CDK6 (PubMed:[19760257](#)). Regulates the expression of HDAC6, ATG7 and VCP in a PPIA/CYPA-dependent manner (PubMed:[25678563](#)).

Cellular Location

Nucleus. Cytoplasm. Cytoplasm, Stress granule Mitochondrion. Note=Continuously travels in and out of the nucleus (PubMed:[18957508](#)). Localizes to stress granules in response to oxidative stress (PubMed:[19765185](#)). A small subset localizes in mitochondria (PubMed:[28794432](#)).

Tissue Location

Ubiquitously expressed. In particular, expression is high in pancreas, placenta, lung, genital tract and spleen

Background


HIV-1, the causative agent of acquired immunodeficiency syndrome (AIDS), contains an RNA genome that produces a chromosomally integrated DNA during the replicative cycle. Activation of HIV-1 gene expression by the transactivator Tat is dependent on an RNA regulatory element (TAR) located downstream of the transcription initiation site. The protein encoded by this gene is a transcriptional repressor that binds to chromosomally integrated TAR DNA and represses HIV-1 transcription. In addition, this protein regulates alternate splicing of the CFTR gene. A similar pseudogene is present on chromosome 20. [provided by RefSeq].

References


- Kim, S.H., et al. J. Biol. Chem. 285(44):34097-34105(2010)
 Geser, F., et al. Arch. Neurol. 67(10):1238-1250(2010)
 Mackenzie, I.R., et al. Lancet Neurol 9(10):995-1007(2010)
 Shan, X., et al. Proc. Natl. Acad. Sci. U.S.A. 107(37):16325-16330(2010)
 McKee, A.C., et al. J. Neuropathol. Exp. Neurol. 69(9):918-929(2010)

Images


Immunohistochemical analysis of AP13763A on paraffin-embedded Human placenta tissue. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:100) for 1 hour at room temperature.



Undiluted CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



All lanes : Anti-TARDBP Antibody (N-term) at 1:2000 dilution Lane 1: HeLa whole cell lysate Lane 2: A431 whole cell lysate Lane 3: HepG2 whole cell lysate Lane 4: K562 whole cell lysate Lane 5: NIH/3T3 whole cell lysate Lane 6: Mouse brain tissue lysate Lane 7: Rat brain tissue lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 45 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Fluorescent confocal image of HeLa cell stained with TARDBP Antibody (N-term)(Cat#AP13763a).HeLa cells were fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.1%, 10 min), then incubated with TARDBP primary antibody (1:25, 1 h at 37°C). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-rabbit antibody (green) was used (1:400, 50 min at 37°C).Cytoplasmic actin was counterstained with Alexa Fluor® 555 (red) conjugated Phalloidin (7units/ml, 1 h at 37°C). Nuclei were counterstained with DAPI (blue) (10 µg/ml, 10 min).TARDBP immunoreactivity is localized to nucleus significantly and Cytoplasm weakly.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.