

Mouse Csk Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP16354b

Product Information

| | |
|--------------------------|---|
| Application | WB, E |
| Primary Accession | P41241 |
| Other Accession | P32577 , P41240 , P41239 , Q0VBZ0 |
| Reactivity | Human, Mouse |
| Predicted | Bovine, Chicken, Rat |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | Rabbit IgG |
| Clone Names | RB30304 |
| Calculated MW | 50716 |
| Antigen Region | 377-404 |

Additional Information

| | |
|---------------------------|--|
| Gene ID | 12988 |
| Other Names | Tyrosine-protein kinase CSK, C-Src kinase, Protein-tyrosine kinase MPK-2, p50CSK, Csk |
| Target/Specificity | This Mouse Csk antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 377-404 amino acids from the C-terminal region of mouse Csk. |
| Dilution | WB~~1:1000 E~~Use at an assay dependent concentration. |
| Format | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| Storage | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| Precautions | Mouse Csk Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures. |

Protein Information

| | |
|-----------------|---|
| Name | Csk |
| Function | Non-receptor tyrosine-protein kinase that plays an important role in the regulation of cell growth, differentiation, migration and immune response. |

Phosphorylates tyrosine residues located in the C-terminal tails of Src-family kinases (SFKs) including LCK, SRC, HCK, FYN, LYN, CSK or YES1. Upon tail phosphorylation, Src-family members engage in intramolecular interactions between the phosphotyrosine tail and the SH2 domain that result in an inactive conformation. To inhibit SFKs, CSK is recruited to the plasma membrane via binding to transmembrane proteins or adapter proteins located near the plasma membrane. Suppresses signaling by various surface receptors, including T-cell receptor (TCR) and B-cell receptor (BCR) by phosphorylating and maintaining inactive several positive effectors such as FYN or LCK (By similarity). May act as a negative regulator of EGFR and STAT3 signaling pathways (By similarity).

Cellular Location

Cytoplasm. Cell membrane. Note=Mainly cytoplasmic, also present in lipid rafts


Tissue Location

Ubiquitous, but most abundant in thymus and spleen, as well as in neonatal brain


Background

Specifically phosphorylates 'Tyr-504' on LCK, which acts as a negative regulatory site. Can also act on the LYN and FYN kinases.

Images



Mouse Csk Antibody (C-term) (Cat. #AP16354b) western blot analysis in Jurkat cell line lysates (35ug/lane). This demonstrates the Csk antibody detected the Csk protein (arrow).



Mouse Csk Antibody (C-term) (Cat. #AP16354b) western blot analysis in mouse spleen tissue lysates (35ug/lane). This demonstrates the Csk antibody detected the Csk protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.