

# OR4D2 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP16376b

#### **Product Information**

**Application** WB, E **Primary Accession** P58180

Other Accession NP\_001004707.1

Reactivity Human
Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Clone Names RB35430
Calculated MW 34958
Antigen Region 211-239

#### **Additional Information**

**Gene ID** 124538

Other Names Olfactory receptor 4D2, B-lymphocyte membrane protein BC2009, Olfactory

receptor OR17-24, OR4D2

**Target/Specificity** This OR4D2 antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 211-239 amino acids from the

C-terminal region of human OR4D2.

**Dilution** WB~~1:1000 E~~Use at an assay dependent concentration.

**Format** Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** OR4D2 Antibody (C-term) is for research use only and not for use in

diagnostic or therapeutic procedures.

#### **Protein Information**

Name OR4D2

**Function** Odorant receptor.

**Cellular Location** Cell membrane; Multi-pass membrane protein.


## **Background**

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

#### References

Zody, M.C., et al. Nature 440(7087):1045-1049(2006) Malnic, B., et al. Proc. Natl. Acad. Sci. U.S.A. 101(8):2584-2589(2004) Fuchs, T., et al. Genomics 80(3):295-302(2002)

### **Images**



OR4D2 Antibody (C-term) (Cat. #AP16376b) western blot analysis in NCI-H460 cell line lysates (35ug/lane). This demonstrates the OR4D2 antibody detected the OR4D2 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.