

Beclin 1 Antibody

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1818a

Product Information

Application	WB, IHC-P, E
Primary Accession	Q14457
Other Accession	Q5ZKS6 , Q4A1L4
Reactivity	Human, Mouse
Predicted	Chicken, Bovine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	51896
Antigen Region	181-210

Additional Information

Gene ID	8678
Other Names	Beclin-1, Coiled-coil myosin-like BCL2-interacting protein, Protein GT197, BECN1, GT197
Target/Specificity	This Beclin 1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 181-210 amino acids from human Beclin 1.
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Beclin 1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	BECN1
Synonyms	GT197
Function	Plays a central role in autophagy (PubMed: 18570871 , PubMed: 21358617 ,

PubMed:[23184933](#), PubMed:[23974797](#), PubMed:[25484083](#), PubMed:[28445460](#), PubMed:[37776275](#)). Acts as a core subunit of the PI3K complex that mediates formation of phosphatidylinositol 3-phosphate; different complex forms are believed to play a role in multiple membrane trafficking pathways: PI3KC3-C1 is involved in initiation of autophagosomes and PI3KC3-C2 in maturation of autophagosomes and endocytosis. Involved in regulation of degradative endocytic trafficking and required for the abscission step in cytokinesis, probably in the context of PI3KC3-C2 (PubMed:[20208530](#), PubMed:[20643123](#), PubMed:[23974797](#), PubMed:[26783301](#)). Essential for the formation of PI3KC3-C2 but not PI3KC3-C1 PI3K complex forms. Involved in endocytosis (PubMed:[25275521](#)). May play a role in antiviral host defense.

Cellular Location

Cytoplasm. Golgi apparatus, trans-Golgi network membrane; Peripheral membrane protein. Endosome membrane; Peripheral membrane protein. Endoplasmic reticulum membrane; Peripheral membrane protein. Mitochondrion membrane; Peripheral membrane protein. Endosome {ECO:0000250|UniProtKB:O88597} Cytoplasmic vesicle, autophagosome. Note=Interaction with ATG14 promotes translocation to autophagosomes. Expressed in dendrites and cell bodies of cerebellar Purkinje cells (By similarity) {ECO:0000250|UniProtKB:O88597, ECO:0000269|PubMed:19050071} [Beclin-1-C 37 kDa]: Mitochondrion {ECO:0000250|UniProtKB:O88597}

Tissue Location

Ubiquitous.

Background


Macroautophagy is the major inducible pathway for the general turnover of cytoplasmic constituents in eukaryotic cells, it is also responsible for the degradation of active cytoplasmic enzymes and organelles during nutrient starvation. Macroautophagy involves the formation of double-membrane bound autophagosomes which enclose the cytoplasmic constituent targeted for degradation in a membrane bound structure, which then fuse with the lysosome (or vacuole) releasing a single-membrane bound autophagic bodies which are then degraded within the lysosome (or vacuole). Beclin 1 plays a role in two fundamentally important cell biological pathways: autophagy and apoptosis. Beclin 1 is thought to function as a VPS and autophagy protein as part of a complex with Class III PI3 kinase, Vps34.

References



Baehrecke EH. *Nat Rev Mol Cell Biol.* 6(6):505-10. (2005) Lum JJ, et al. *Nat Rev Mol Cell Biol.* 6(6):439-48. (2005) Greenberg JT. *Dev Cell.* 8(6):799-801. (2005) Levine B. *Cell.* 120(2):159-62. (2005) Shintani T and Klionsky DJ. *Science.* 306(5698):990-5. (2004) Liang X.H., et al. *J. Virol.* 72 (11), 8586-8596 (1998) Aita V.M., et al. *Genomics* 59:59-65(1999).

Images

Western blot analysis of anti-hBECN1-Q196 Pab (Cat. #AP1818a) in 293 cell line lysates transiently transfected with the BECN1 gene (2ug/lane). hBECN1-Q196(arrow) was detected using the purified Pab.



The anti-BECN1 Pab (Cat. #AP1818a) is used in Western blot to detect BECN1 in mouse liver tissue lysate. BECN1 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

Citations

- [Expression of LC3 and Beclin 1 in the spinal dorsal horn following spinal nerve ligation-induced neuropathic pain.](#)
- [Triterpenes from Ganoderma Lucidum induce autophagy in colon cancer through the inhibition of p38 mitogen-activated kinase \(p38 MAPK\).](#)
- [Chondrocyte autophagy is stimulated by HIF-1 dependent AMPK activation and mTOR suppression.](#)
- [Down-regulation of autophagy-related protein 5 \(ATG5\) contributes to the pathogenesis of early-stage cutaneous melanoma.](#)
- [TLR4-mediated autophagic impairment contributes to neuropathic pain in chronic constriction injury mice.](#)
- [Curcumin induces apoptotic cell death and protective autophagy in human gastric cancer cells.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.