

# GTPBP3 Antibody(Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP19547c

#### **Product Information**

**Application** WB, E **Primary Accession Q969Y2** Other Accession NP 116009.2 Reactivity Human Host Rabbit Clonality Polyclonal Isotype Rabbit IgG **Clone Names** RB40710 **Calculated MW** 52058 293-320 **Antigen Region** 

### **Additional Information**

**Gene ID** 84705

Other Names tRNA modification GTPase GTPBP3, mitochondrial, GTP-binding protein 3,

Mitochondrial GTP-binding protein 1, GTPBP3, MTGP1

**Target/Specificity**This GTPBP3 antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 293-320 amino acids from the Central

region of human GTPBP3.

**Dilution** WB~~1:1000 E~~Use at an assay dependent concentration.

**Format** Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** GTPBP3 Antibody(Center) is for research use only and not for use in

diagnostic or therapeutic procedures.

#### **Protein Information**

Name GTPBP3 ( HGNC:14880)

Synonyms MTGP1

**Function** GTPase component of the GTPBP3-MTO1 complex that catalyzes the

5-taurinomethyluridine (taum(5)U) modification at the 34th wobble position (U34) of mitochondrial tRNAs (mt-tRNAs), which plays a role in mt-tRNA decoding and mitochondrial translation (PubMed:29390138, PubMed:33619562). Taum(5)U formation on mammalian mt-tRNA requires the presence of both GTPBP3-mediated GTPase activity and MTO1 catalytic activity (PubMed:29390138).

Cellular Location Mitochondrion

**Tissue Location** Ubiquitously expressed.


## **Background**

This locus encodes a GTP-binding protein. The encoded protein is localized to the mitochondria and may play a role in mitochondrial tRNA modification. Polymorphisms at this locus may be associated with severity of aminoglycoside-induced deafness, a disease associated with a mutation in the 12S rRNA. Alternatively spliced transcript variants encoding different isoforms have been described.

#### References

Reiling, E., et al. Eur. J. Hum. Genet. 17(8):1056-1062(2009) Villarroya, M., et al. Mol. Cell. Biol. 28(24):7514-7531(2008) Bykhovskaya, Y., et al. Mol. Genet. Metab. 83(3):199-206(2004) Li, X., et al. Biochem. Biophys. Res. Commun. 312(3):747-754(2003) Li, X., et al. Mol. Cell. Biol. 22(21):7701-7711(2002)

## **Images**



GTPBP3 Antibody (Center) (Cat. #AP19547c) western blot analysis in NCI-H460 cell line lysates (35ug/lane). This demonstrates the GTPBP3 antibody detected the GTPBP3 protein (arrow).

 $Please\ note: All\ products\ are\ 'FOR\ RESEARCH\ USE\ ONLY.\ NOT\ FOR\ USE\ IN\ DIAGNOSTIC\ OR\ THERAPEUTIC\ PROCEDURES'.$