

UNC5C Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP19910a

Product Information

Application	WB, E
Primary Accession	O95185
Other Accession	NP_003719.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB41789
Calculated MW	103146
Antigen Region	88-117

Additional Information

Gene ID	8633
Other Names	Netrin receptor UNC5C, Protein unc-5 homolog 3, Protein unc-5 homolog C, UNC5C, UNC5H3
Target/Specificity	This UNC5C antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 88-117 amino acids from the N-terminal region of human UNC5C.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	UNC5C Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	UNC5C
Synonyms	UNC5H3
Function	Receptor for netrin required for axon guidance (By similarity). Mediates

axon repulsion of neuronal growth cones in the developing nervous system upon ligand binding (By similarity). NTN1/Netrin-1 binding might cause dissociation of UNC5C from polymerized TUBB3 in microtubules and thereby lead to increased microtubule dynamics and axon repulsion (PubMed:[28483977](#)). Axon repulsion in growth cones may also be caused by its association with DCC that may trigger signaling for repulsion (By similarity). Might also collaborate with DSCAM in NTN1-mediated axon repulsion independently of DCC (By similarity). Also involved in corticospinal tract axon guidance independently of DCC (By similarity). Involved in dorsal root ganglion axon projection towards the spinal cord (PubMed:[28483977](#)). It also acts as a dependence receptor required for apoptosis induction when not associated with netrin ligand (By similarity).

Cellular Location

Cell membrane; Single-pass type I membrane protein. Cell surface. Synapse, synaptosome {ECO:0000250|UniProtKB:Q761X5}. Cell projection, axon {ECO:0000250|UniProtKB:O08747}. Cell projection, dendrite {ECO:0000250|UniProtKB:O08747}. Cell projection, growth cone {ECO:0000250|UniProtKB:O08747}. Cell projection, lamellipodium {ECO:0000250|UniProtKB:O08747}. Cell projection, filopodium {ECO:0000250|UniProtKB:O08747}


Tissue Location

Mainly expressed in brain (PubMed:9782087). Expressed in temporal lobe cortical neurons and in neurons of the hippocampal pyramidal layer (PubMed:25419706). Also expressed in kidney (PubMed:9782087). Not expressed in developing or adult lung (PubMed:9782087).

Background


Receptor for netrin required for axon guidance. Mediates axon repulsion of neuronal growth cones in the developing nervous system upon ligand binding. Axon repulsion in growth cones may be caused by its association with DCC that may trigger signaling for repulsion. Also involved in corticospinal tract axon guidances independently of DCC. It also acts as a dependence receptor required for apoptosis induction when not associated with netrin ligand (By similarity).

Images



Anti-UNC5C Antibody (N-term) at 1:1000 dilution + Jurkat whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 103 kDa Blocking/Dilution buffer: 5% NFDm/TBST.

UNC5C Antibody (N-term) (Cat. #AP19910a) western blot analysis in CEM cell line lysates (35ug/lane). This demonstrates the UNC5C antibody detected the UNC5C protein (arrow).



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.