

CLDN14 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP20035b

Product Information

| | |
|--------------------------|-----------------------------|
| Application | WB, E |
| Primary Accession | O95500 |
| Other Accession | NP_036262.1 |
| Reactivity | Human, Mouse |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | Rabbit IgG |
| Clone Names | RB42184 |
| Calculated MW | 25699 |
| Antigen Region | 172-200 |

Additional Information

| | |
|---------------------------|--|
| Gene ID | 23562 |
| Other Names | Claudin-14, CLDN14 |
| Target/Specificity | This CLDN14 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 172-200 amino acids from the C-terminal region of human CLDN14. |
| Dilution | WB~~1:1000 E~~Use at an assay dependent concentration. |
| Format | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| Storage | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| Precautions | CLDN14 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures. |

Protein Information

| | |
|--------------------------|--|
| Name | CLDN14 |
| Function | Plays a major role in tight junction-specific obliteration of the intercellular space, through calcium-independent cell-adhesion activity. |
| Cellular Location | Cell junction, tight junction. Cell membrane; Multi-pass membrane protein |

Tissue Location

Liver, kidney. Also found in ear.


Background

Tight junctions represent one mode of cell-to-cell adhesion in epithelial or endothelial cell sheets, forming continuous seals around cells and serving as a physical barrier to prevent solutes and water from passing freely through the paracellular space. These junctions are comprised of sets of continuous networking strands in the outwardly facing cytoplasmic leaflet, with complementary grooves in the inwardly facing extracytoplasmic leaflet. The protein encoded by this gene, a member of the claudin family, is an integral membrane protein and a component of tight junction strands. The encoded protein also binds specifically to the WW domain of Yes-associated protein. Defects in this gene are the cause of an autosomal recessive form of nonsyndromic sensorineural deafness. It is also reported that four synonymous variants in this gene are associated with kidney stones and reduced bone mineral density. Several transcript variants encoding the same protein have been found for this gene. [provided by RefSeq].


References

Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :
Thorleifsson, G., et al. Nat. Genet. 41(8):926-930(2009)
Belguith, H., et al. Biochem. Biophys. Res. Commun. 385(1):1-5(2009)
Lal-Nag, M., et al. Genome Biol. 10 (8), 235 (2009) :
Krause, G., et al. Biochim. Biophys. Acta 1778(3):631-645(2008)

Images



Western blot analysis of lysates from human kidney and liver tissue lysate (from left to right), using CLDN14 Antibody (C-term) (Cat. #AP20035b). AP20035b was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35ug per lane.



CLDN14 Antibody (C-term) (Cat. #AP20035b) western blot analysis in MDA-MB453 cell line lysates (35ug/lane). This demonstrates the CLDN14 antibody detected the CLDN14 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.