

BMPR1B Antibody (N-term K15)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP2005a

Product Information

Application	WB, E
Primary Accession	O00238
Other Accession	NP_001194
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB1774
Calculated MW	56930
Antigen Region	1-30

Additional Information

Gene ID	658
Other Names	Bone morphogenetic protein receptor type-1B, BMP type-1B receptor, BMPR-1B, CDw293, BMPR1B
Target/Specificity	This BMPR1B antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human BMPR1B.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	BMPR1B Antibody (N-term K15) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	BMPR1B
Function	On ligand binding, forms a receptor complex consisting of two type II and two type I transmembrane serine/threonine kinases. Type II receptors phosphorylate and activate type I receptors which autophosphorylate, then

bind and activate SMAD transcriptional regulators. Receptor for BMP7/OP-1 and GDF5. Positively regulates chondrocyte differentiation through GDF5 interaction.

Cellular Location

Cell membrane; Single-pass type I membrane protein


Background

The bone morphogenetic protein (BMP) receptors are a family of transmembrane serine/threonine kinases that include the type I receptors BMPR1A and BMPR1B and the type II receptor BMPR2. These receptors are also closely related to the activin receptors, ACVR1 and ACVR2. The ligands of these receptors are members of the TGF-beta superfamily. TGF-betas and activins transduce their signals through the formation of heteromeric complexes with 2 different types of serine (threonine) kinase receptors: type I receptors of about 50-55 kD and type II receptors of about 70-80 kD. Type II receptors bind ligands in the absence of type I receptors, but they require their respective type I receptors for signaling, whereas type I receptors require their respective type II receptors for ligand binding.

References

- Lehmann, K., et al., Proc. Natl. Acad. Sci. U.S.A. 100(21):12277-12282 (2003).
Astrom, A.K., et al., Mamm. Genome 10(3):299-302 (1999).
Ide, H., et al., Oncogene 14(11):1377-1382 (1997).
ten Dijke, P., et al., Science 264(5155):101-104 (1994).
Ide, H., et al., Cytogenet. Cell Genet. 81 (3-4), 285-286 (1998).

Images



Western blot analysis of anti-BMPR1B Pab (Cat. #ap2005a) in NCI-H460 cell lysate. BMPR1B (arrow) was detected using purified Pab. Secondary HRP-anti-rabbit was used for signal visualization with chemiluminescence.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.