

EXOC4 Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21329a

Product Information

Application	WB, E
Primary Accession	Q96A65
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB52634
Calculated MW	110498

Additional Information

Gene ID	60412
Other Names	Exocyst complex component 4, Exocyst complex component Sec8, EXOC4, KIAA1699, SEC8, SEC8L1
Target/Specificity	This EXOC4 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 239-273 amino acids from the N-terminal region of human EXOC4.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	EXOC4 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	EXOC4
Synonyms	KIAA1699, SEC8, SEC8L1
Function	Component of the exocyst complex involved in the docking of exocytic vesicles with fusion sites on the plasma membrane.

Cellular Location

Midbody, Midbody ring. Cell projection {ECO:0000250|UniProtKB:Q62824}. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome Note=Colocalizes with CNTRL/centriolin at the midbody ring (PubMed:16213214). Localizes at the leading edge of migrating cells (By similarity). {ECO:0000250|UniProtKB:Q62824, ECO:0000269|PubMed:16213214}

Background

Component of the exocyst complex involved in the docking of exocytic vesicles with fusion sites on the plasma membrane.

References

Sha J.H.,et al.Submitted (MAY-2001) to the EMBL/GenBank/DDBJ databases.


Ota T.,et al.Nat. Genet. 36:40-45(2004).

Bechtel S.,et al.BMC Genomics 8:399-399(2007).

Scherer S.W.,et al.Science 300:767-772(2003).

Hillier L.W.,et al.Nature 424:157-164(2003).

Images



All lanes : Anti-EXOC4 Antibody (N-term) at 1:2000 dilution Lane 1: HeLa whole cell lysates Lane 2: Jurkat whole cell lysates Lane 3: MCF-7 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 110 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.