

TNFRSF14 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21380b

Product Information

| | |
|--------------------------|------------------------|
| Application | WB, E |
| Primary Accession | Q92956 |
| Reactivity | Human |
| Host | Rabbit |
| Clonality | polyclonal |
| Isotype | Rabbit IgG |
| Clone Names | RB52847 |

Additional Information

| | |
|---------------------------|--|
| Other Names | Tumor necrosis factor receptor superfamily member 14, Herpes virus entry mediator A, Herpesvirus entry mediator A, HveA, Tumor necrosis factor receptor-like 2, TR2, CD270, TNFRSF14, HVEA, HVEM |
| Target/Specificity | This TNFRSF14 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 269-302 amino acids from the C-terminal region of human TNFRSF14. |
| Dilution | WB~~1:1000 E~~Use at an assay dependent concentration. |
| Format | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| Storage | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| Precautions | TNFRSF14 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures. |

Protein Information

Background

Receptor for BTLA. Receptor for TNFSF14/LIGHT and homotrimeric TNFSF1/lymphotoxin-alpha. Involved in lymphocyte activation. Plays an important role in HSV pathogenesis because it enhanced the entry of several wild-type HSV strains of both serotypes into CHO cells, and mediated HSV entry into activated human T-cells.

References

Montgomery R.I.,et al.Cell 87:427-436(1996).


Kwon B.S.,et al.J. Biol. Chem. 272:14272-14276(1997).

Zhang W.,et al.Submitted (MAY-1999) to the EMBL/GenBank/DDBJ databases.

Struyf F.,et al.J. Infect. Dis. 185:36-44(2002).

Ota T.,et al.Nat. Genet. 36:40-45(2004).

Images



Anti-TNFRSF14 Antibody (C-term)at 1:1000 dilution + Raji whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 30 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.