

Mouse Cxxc1 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21466c

Product Information

Application	WB, E
Primary Accession	Q9CWW7
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB50989
Calculated MW	76167

Additional Information

Gene ID	74322
Other Names	CXXC-type zinc finger protein 1, CpG-binding protein, PHD finger and CXXC domain-containing protein 1, Cxxc1, Cgbp, Pccx1
Target/Specificity	This Mouse Cxxc1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 364-399 amino acids from the Central region of Mouse Cxxc1.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Mouse Cxxc1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	Cxxc1
Synonyms	Cgbp, Pccx1
Function	Transcriptional activator that exhibits a unique DNA binding specificity for CpG unmethylated motifs with a preference for CpGG.

Cellular Location	Nucleus speckle {ECO:0000250 UniProtKB:Q9P0U4}. Nucleus. Note=Associated with euchromatin. During mitosis, excluded from condensed chromosomes (By similarity). {ECO:0000250 UniProtKB:Q9P0U4}
Tissue Location	Expressed in seminiferous tubules and in both germ cells and Sertoli cells. Highly expressed in spermatogonia, weakly expressed in leptoneuma and zygonema, and then again high expression in pachynema and diplonema, decreasing to undetectable levels in spermatids.


Background

Transcriptional activator that exhibits a unique DNA binding specificity for CpG unmethylated motifs with a preference for CpGG.

References

Carninci P., et al. Science 309:1559-1563(2005).

Images



Anti-Cxxc1 Antibody (Center) at 1:2000 dilution + mouse thymus lysates. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 76 kDa. Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.