

SRY Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21539c

Product Information

Application	WB, E
Primary Accession	Q05066
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB53826
Calculated MW	23884

Additional Information

Gene ID	6736
Other Names	Sex-determining region Y protein, Testis-determining factor, SRY, TDF
Target/Specificity	This SRY antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 81-112 amino acids from the Central region of human SRY.
Dilution	WB~1:1000 E~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	SRY Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	SRY {ECO:0000303 PubMed:1695712, ECO:0000312 HGNC:HGNC:11311}
Function	Transcriptional regulator that controls a genetic switch in male development (PubMed: 11563911). It is necessary and sufficient for initiating male sex determination by directing the development of supporting cell precursors (pre-Sertoli cells) as Sertoli rather than granulosa cells (PubMed: 16414182 , PubMed: 16996051). Involved in different aspects of gene regulation including promoter activation or repression (PubMed: 9525897).

Binds to the DNA consensus sequence 5'- [AT]AACAA[AT]-3' (PubMed:[11563911](#), PubMed:[1425584](#), PubMed:[15170344](#), PubMed:[8159753](#), PubMed:[8265659](#)). SRY HMG box recognizes DNA by partial intercalation in the minor groove and promotes DNA bending (PubMed:[11563911](#), PubMed:[1425584](#), PubMed:[15170344](#), PubMed:[16762365](#), PubMed:[8159753](#), PubMed:[8265659](#)). Also involved in pre-mRNA splicing (PubMed:[11818535](#)). In male adult brain involved in the maintenance of motor functions of dopaminergic neurons (By similarity).

Cellular Location

Nucleus speckle. Cytoplasm Nucleus. Note=Acetylation contributes to its nuclear localization and deacetylation by HDAC3 induces a cytoplasmic delocalization (PubMed:15297880). Colocalizes with SOX6 in speckles (PubMed:11818535). Colocalizes with CAML in the nucleus (PubMed:15746192). Colocalizes in the nucleus with ZNF208 isoform KRAB- O and tyrosine hydroxylase (TH) (By similarity). Nuclear import is facilitated by XPO4, a protein that usually acts as a nuclear export signal receptor (PubMed:19349578). {ECO:0000250|UniProtKB:Q05738, ECO:0000269|PubMed:11818535, ECO:0000269|PubMed:15297880, ECO:0000269|PubMed:15746192, ECO:0000269|PubMed:19349578}


Background

Transcriptional regulator that controls a genetic switch in male development. It is necessary and sufficient for initiating male sex determination by directing the development of supporting cell precursors (pre-Sertoli cells) as Sertoli rather than granulosa cells (By similarity). In male adult brain involved in the maintenance of motor functions of dopaminergic neurons (By similarity). Involved in different aspects of gene regulation including promoter activation or repression (By similarity). Promotes DNA bending. SRY HMG box recognizes DNA by partial intercalation in the minor groove. Also involved in pre-mRNA splicing. Binds to the DNA consensus sequence 5'-[AT]AACAA[AT]-3'.

References

- Sinclair A.H.,et al.Nature 346:240-244(1990).
Su H.,et al.Am. J. Hum. Genet. 52:24-38(1993).
Behlke M.A.,et al.Genomics 17:736-739(1993).
Whitfield L.S.,et al.Genomics 27:306-311(1995).
Ferrari S.,et al.EMBO J. 11:4497-4506(1992).

Images



Anti-SRY Antibody (Center)at 1:1000 dilution + K562 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 24 kDa Blocking/Dilution buffer: 5% NFDM/TBST.